

## PRESCRIBING INFORMATION

### Pneumococcal 7-valent Conjugate Vaccine (Diphtheria CRM<sub>197</sub> Protein)

*Pprevnar*<sup>™</sup>

Rx only

#### ***For Intramuscular Injection Only***

#### **DESCRIPTION**

Pprevnar<sup>™</sup>, Pneumococcal 7-valent Conjugate Vaccine (Diphtheria CRM<sub>197</sub> Protein), is a sterile solution of saccharides of the capsular antigens of *Streptococcus pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F individually conjugated to diphtheria CRM<sub>197</sub> protein. Each serotype is grown in soy peptone broth. The individual polysaccharides are purified through centrifugation, precipitation, ultrafiltration, and column chromatography. The polysaccharides are chemically activated to make saccharides which are directly conjugated to the protein carrier CRM<sub>197</sub> to form the glycoconjugate. This is effected by reductive amination. CRM<sub>197</sub> is a nontoxic variant of diphtheria toxin isolated from cultures of *Corynebacterium diphtheriae* strain C7 (β197) grown in a casamino acids and yeast extract-based medium. CRM<sub>197</sub> is purified through ultrafiltration, ammonium sulfate precipitation, and ion-exchange chromatography. The individual glycoconjugates are purified by ultrafiltration and column chromatography and are analyzed for saccharide to protein ratios, molecular size, free saccharide, and free protein.

The individual glycoconjugates are compounded to formulate the vaccine, Pprevnar<sup>™</sup>. Potency of the formulated vaccine is determined by quantification of each of the saccharide antigens, and by the saccharide to protein ratios in the individual glycoconjugates.

Pprevnar<sup>™</sup> is manufactured as a liquid preparation. Each 0.5 mL dose is formulated to contain: 2 µg of each saccharide for serotypes 4, 9V, 14, 18C, 19F, and 23F, and 4 µg of serotype 6B per dose (16 µg total saccharide); approximately 20 µg of CRM<sub>197</sub> carrier protein; and 0.125 mg of aluminum per 0.5 mL dose as aluminum phosphate adjuvant.

After shaking, the vaccine is a homogeneous, white suspension.

#### **CLINICAL PHARMACOLOGY**

*S. pneumoniae* is an important cause of morbidity and mortality in persons of all ages worldwide. The organism causes invasive infections, such as bacteremia and meningitis, as well as pneumonia and upper respiratory tract infections including otitis media and sinusitis. In children older than 1 month, *S. pneumoniae* is the most common cause of invasive disease.<sup>1</sup> Data from community-based studies performed between 1986 and 1995, indicate that the overall annual incidence of invasive pneumococcal disease in the United States is an estimated 10 to 30 cases per 100,000 persons, with the highest risk in children aged less than or equal to 2 years of age (140 to 160 cases per 100,000 persons).<sup>2,3,4,5,6</sup> Children in group child care have an increased risk for invasive pneumococcal disease.<sup>7,8</sup> Immunocompromised individuals with neutropenia, asplenia, sickle cell disease, disorders of complement and humoral immunity, human immunodeficiency virus (HIV) infections or chronic underlying disease are also at increased risk for invasive pneumococcal disease.<sup>8</sup> *S. pneumoniae* is the most common cause of bacterial meningitis in the United States.<sup>1</sup> The annual incidence of pneumococcal meningitis in children between 1 to 23 months of age is approximately 7 cases per 100,000 persons.<sup>1</sup> Pneumococcal meningitis in childhood has been associated with 8% mortality and may result in neurological sequelae (25%) and hearing loss (32%) in survivors.<sup>9</sup>

*S. pneumoniae* is an important cause of acute otitis media, identified in 20 to 40% of middle ear fluid cultures.<sup>10,11</sup> The seven serotypes account for approximately 60% of acute otitis media due to *S. pneumoniae* (12-24% of all acute otitis media).<sup>12</sup> The exact contribution of *S. pneumoniae* to childhood pneumonia is unknown, as it is often not possible to identify the causative organisms. In studies of children less than 5 years of age with community-acquired pneumonia, where diagnosis

was attempted using serological methods, antigen testing, or culture data, 30% of cases were classified as bacterial pneumonia, and 70% of these (21% of total community-acquired pneumonia) were found to be due to *S. pneumoniae*.<sup>13,14</sup>

In the past decade the proportion of *S. pneumoniae* isolates resistant to antibiotics has been on the rise in the United States and worldwide. In a multi-center US surveillance study, the prevalence of penicillin and cephalosporin-nonsusceptible (intermediate or high level resistance) invasive disease isolates from children was 21% (range < 5% to 38% among centers), and 9.3% (range 0-18%), respectively. Over the 3-year surveillance period (1993-1996), there was a 50% increase in penicillin-nonsusceptible *S. pneumoniae* (PNSP) strains and a three-fold rise in cephalosporin-nonsusceptible strains.<sup>8</sup> Although generally less common than PNSP, pneumococci resistant to macrolides and trimethoprim-sulfazoxole have also been observed.<sup>4</sup> Day care attendance, a history of ear infection, and a recent history of antibiotic exposure, have also been associated with invasive infections with PNSP in children 2 months to 59 months of age.<sup>7,8</sup> There has been no difference in mortality associated with PNSP strains.<sup>8,9</sup> However, the American Academy of Pediatrics (AAP) revised the antibiotic treatment guidelines in 1997 in response to the increased prevalence of antibiotic-resistant pneumococci.<sup>15</sup>

Approximately 90 serotypes of *S. pneumoniae* have been identified based on antigenic differences in their capsular polysaccharides. The distribution of serotypes responsible for disease differ with age and geographic location.<sup>16</sup>

Serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F have been responsible for approximately 80% of invasive pneumococcal disease in children < 6 years of age in the United States.<sup>12</sup> These 7 serotypes also accounted for 74% of PNSP and 100% of pneumococci with high level penicillin resistance isolated from children < 6 years with invasive disease during a 1993-1994 surveillance by the Centers for Disease Control.<sup>17</sup>

## **Results of Clinical Evaluations**

### **Efficacy**

Efficacy was assessed in a randomized, double-blinded clinical trial in a multiethnic population at Northern California Kaiser Permanente (NCKP), beginning in October 1995, in which 37,816 infants were randomized to receive either Prevnar™ or a control vaccine (an investigational meningococcal group C conjugate vaccine [MnCC]) at 2, 4, 6, and 12-15 months of age. Prevnar™ was administered to 18,906 children and the control vaccine to 18,910 children. Routinely recommended vaccines were also administered which changed during the trial to reflect changing AAP and Advisory Committee on Immunization Practices (ACIP) recommendations. A planned interim analysis was performed upon accrual of 17 cases of invasive disease due to vaccine-type *S. pneumoniae* (August 1998). Ancillary endpoints for evaluation of efficacy against pneumococcal disease were also assessed in this trial.

Efficacy against invasive disease: Invasive disease was defined as isolation and identification of *S. pneumoniae* from normally sterile body sites in children presenting with an acute illness consistent with pneumococcal disease. Weekly surveillance of listings of cultures from the NCKP Regional Microbiology database was conducted to assure ascertainment of all cases. The primary endpoint was efficacy against invasive pneumococcal disease due to vaccine serotypes. The per protocol analysis of the primary endpoint included cases which occurred  $\geq 14$  days after the third dose. The intent-to-treat (ITT) analysis included all cases of invasive pneumococcal disease due to vaccine serotypes in children who received at least one dose of vaccine. Secondary analyses of efficacy against all invasive pneumococcal disease, regardless of serotype, were also performed according to these same per protocol and ITT definitions. Results of these analyses are presented in Table 1.

**TABLE 1**  
**Efficacy of Prevnar™ Against Invasive Disease Due to *S. pneumoniae***  
**in Cases Accrued From October 15, 1995 Through August 20, 1998<sup>18,19</sup>**

	Prevnar™	Control*	Efficacy	95% CI
	Number of Cases	Number of Cases		
<b>Vaccine serotypes</b>				
<b>Per protocol</b>	<b>0</b>	<b>17</b>	<b>100%</b>	<b>75.4, 100</b>
<b>Intent-to-treat</b>	<b>0</b>	<b>22</b>	<b>100%</b>	<b>81.7, 100</b>
<b>All pneumococcal serotypes</b>				
Per protocol	2	20	90.0%	58.3, 98.9
Intent-to-treat	3	27†	88.9%	63.8, 97.9

\* Investigational meningococcal group C conjugate vaccine (MnCC).

† Includes one case in an immunocompromised subject.

All 22 cases of invasive disease due to vaccine serotype strains in the ITT population were bacteremic. In addition, the following diagnoses were also reported: meningitis (2), pneumonia (2), and cellulitis (1).

Preliminary efficacy data through an extended follow-up period to April 20, 1999, resulted in a similar efficacy estimate (Per protocol: 1 case in Prevnar™ group, 39 cases in control group; ITT: 3 cases in Prevnar™ group, 49 cases in the control group).

### **Immunogenicity**

#### **Routine Schedule**

Subjects from a subset of selected study sites in the NCKP efficacy study were approached for participation in the immunogenicity portion of the study on a volunteer basis. Immune responses following three or four doses of Prevnar™ or the control vaccine were evaluated in children who received either concurrent Diphtheria and Tetanus Toxoids and Pertussis Vaccine Adsorbed and Haemophilus b Conjugate Vaccine (Diphtheria CRM<sub>197</sub> Protein Conjugate), (DTP-HbOC), or Diphtheria and Tetanus Toxoids and Acellular Pertussis Vaccine Adsorbed (DTaP), and Haemophilus b Conjugate Vaccine (Diphtheria CRM<sub>197</sub> Protein Conjugate), (HbOC) vaccines at 2, 4, and 6 months of age. The use of Hepatitis B (Hep B), Oral Polio Vaccine (OPV), Inactivated Polio Vaccine (IPV), Measles-Mumps-Rubella (MMR), and Varicella vaccines were permitted according to the AAP and ACIP recommendations.

Table 2 presents the geometric mean concentrations (GMC) of pneumococcal antibodies following the third and fourth doses of Prevnar™ or the control vaccine when administered concurrently with DTP-HbOC vaccine in the efficacy study.

**TABLE 2**  
**Geometric Mean Concentrations (µg/mL) of Pneumococcal Antibodies Following the Third and Fourth Doses of Prevnar™ or Control\* When Administered Concurrently With DTP-HbOC in the Efficacy Study<sup>19</sup>**

Serotype	Post dose 3 GMC† (95% CI for Prevnar™)		Post dose 4 GMC‡ (95% CI for Prevnar™)	
	Prevnar™§	Control*	Prevnar™§	Control*
	N=88	N=92	N=68	N=61
4	1.46 (1.19, 1.78)	0.03	2.38 (1.88, 3.03)	0.04
6B	4.70 (3.59, 6.14)	0.08	14.45 (11.17, 18.69)	0.17
9V	1.99 (1.64, 2.42)	0.05	3.51 (2.75, 4.48)	0.06
14	4.60 (3.70, 5.74)	0.05	6.52 (5.18, 8.21)	0.06
18C	2.16 (1.73, 2.69)	0.04	3.43 (2.70, 4.37)	0.07
19F	1.39 (1.16, 1.68)	0.09	2.07 (1.66, 2.57)	0.18
23F	1.85 (1.46, 2.34)	0.05	3.82 (2.85, 5.11)	0.09

\* Control was investigational meningococcal group C conjugate vaccine (MnCC).

† Mean age of Prevnar™ group was 7.8 months and of control group was 7.7 months.

N is slightly less for some serotypes in each group.

‡ Mean age of Prevnar™ group was 14.2 months and of control group was 14.4 months.

N is slightly less for some serotypes in each group.

§ p<0.001 when Prevnar™ compared to control for each serotype using a Wilcoxon's test.

In another randomized study (Manufacturing Bridging Study, 118-16), immune responses were evaluated following three doses of Prevnar™ administered concomitantly with DTaP and HbOC vaccines at 2, 4, and 6 months of age, IPV at 2 and 4 months of age, and Hep B at 2 and 6 months of age. The control group received concomitant vaccines only. Table 3 presents the immune responses to pneumococcal polysaccharides observed in both this study and in the subset of subjects from the efficacy study that received concomitant DTaP and HbOC vaccines.

**TABLE 3**  
**Geometric Mean Concentrations (µg/mL) of Pneumococcal Antibodies Following the Third Dose of Prevnar™ or Control\* When Administered Concurrently With DTaP and HbOC in the Efficacy Study† and Manufacturing Bridging Study<sup>19,20</sup>**

Serotype	Efficacy Study		Manufacturing Bridging Study	
	Post dose 3 GMC‡ (95% CI for Prevnar™)		Post dose 3 GMC§ (95% CI for Prevnar™)	
	Prevnar™II	Control*	Prevnar™II	Control*
	N=32	N=32	N=159	N=83
4	1.47 (1.08, 2.02)	0.02	2.03 (1.75, 2.37)	0.02
6B	2.18 (1.20, 3.96)	0.06	2.97 (2.43, 3.65)	0.07
9V	1.52 (1.04, 2.22)	0.04	1.18 (1.01, 1.39)	0.04
14	5.05 (3.32, 7.70)	0.04	4.64 (3.80, 5.66)	0.04
18C	2.24 (1.65, 3.02)	0.04	1.96 (1.66, 2.30)	0.04
19F	1.54 (1.09, 2.17)	0.10	1.91 (1.63, 2.25)	0.08
23F	1.48 (0.97, 2.25)	0.05	1.71 (1.44, 2.05)	0.05

\* Control in efficacy study was investigational meningococcal group C conjugate vaccine (MnCC) and in Manufacturing Bridging Study was concomitant vaccines only.

† Sufficient data are not available to reliably assess GMCs following 4 doses of Prevnar™ when administered with DTaP in the NCKP efficacy study.

‡ Mean age of the Prevnar™ group was 7.4 months and of the control group was 7.6 months. N is slightly less for some serotypes in each group.

§ Mean age of the Prevnar™ group and the control group was 7.2 months.

||  $p < 0.001$  when Prevnar™ compared to control for each serotype using a Wilcoxon's test in the efficacy study and two-sample t-test in the Manufacturing Bridging Study.

In all studies in which the immune responses to Prevnar™ were contrasted to control, a significant antibody response was seen to all vaccine serotypes following three or four doses, although geometric mean concentrations of antibody varied among serotypes.<sup>18,19,20,21,22,23,24,25</sup> The minimum

serum antibody concentration necessary for protection against invasive pneumococcal disease has not been determined for any serotype.

Prevnar™ induces functional antibodies to all vaccine serotypes, as measured by opsonophagocytosis following three doses.<sup>25</sup>

### Previously Unvaccinated Older Infants and Children

To determine an appropriate schedule for children 7 months of age or older at the time of the first immunization with Pevnar™, 483 children in 4 ancillary studies received Pevnar™ at various schedules. GMCs attained using the various schedules among older infants and children were comparable to immune responses of children, who received concomitant DTaP, in the NCKP efficacy study (118-8) after 3 doses for most serotypes, as shown in Table 4. These data support the schedule for previously unvaccinated older infants and children who are beyond the age of the infant schedule. For usage in older infants and children see DOSAGE AND ADMINISTRATION.

**TABLE 4**  
**Geometric Mean Concentrations (µg/mL) of Pneumococcal Antibodies Following Immunization of Children From 7 Months Through 9 Years of Age With Pevnar™<sup>26</sup>**

Age group, Vaccinations	Study	Sample Size(s)	4	6B	9V	14	18C	19F	23F
7-11 mo. 3 doses	118-12	22	<b>2.34</b>	<b>3.66</b>	<b>2.11</b>	<b>9.33</b>	<b>2.31</b>	<b>1.60</b>	<b>2.50</b>
	118-16	39	<b>3.60</b>	<b>4.63</b>	<b>2.04</b>	<b>5.48</b>	<b>1.98</b>	<b>2.15</b>	<b>1.93</b>
12-17 mo. 2 doses	118-15*	82-84†	<b>3.91</b>	<b>4.67</b>	<b>1.94</b>	<b>6.92</b>	<b>2.25</b>	<b>3.78</b>	<b>3.29</b>
	118-18	33	<b>7.02</b>	<b>4.25</b>	<b>3.26</b>	<b>6.31</b>	<b>3.60</b>	<b>3.29</b>	<b>2.92</b>
18-23 mo. 2 doses	118-15*	52-54†	<b>3.36</b>	<b>4.92</b>	<b>1.80</b>	<b>6.69</b>	<b>2.65</b>	<b>3.17</b>	<b>2.71</b>
	118-18	45	<b>6.85</b>	<b>3.71</b>	<b>3.86</b>	<b>6.48</b>	<b>3.42</b>	<b>3.86</b>	<b>2.75</b>
24-35 mo. 1 dose	118-18	53	<b>5.34</b>	<b>2.90</b>	<b>3.43</b>	1.88	<b>3.03</b>	<b>4.07</b>	<b>1.56</b>
36-59 mo. 1 dose	118-18	52	<b>6.27</b>	<b>6.40</b>	<b>4.62</b>	<b>5.95</b>	<b>4.08</b>	<b>6.37</b>	<b>2.95</b>
5-9 yrs. 1 dose	118-18	101	<b>6.92</b>	<b>20.84</b>	<b>7.49</b>	<b>19.32</b>	<b>6.72</b>	<b>12.51</b>	<b>11.57</b>
118-8, DTaP	Post dose 3	31-32†	1.47	2.18	1.52	5.05	2.24	1.54	1.48

Bold = GMC not inferior to 118-8, DTaP post dose 3 (one-sided lower limit of the 95% CI of GMC ratio  $\geq 0.50$ ).

\* Study in Navajo and Apache populations.

† Numbers vary with serotype.

### INDICATIONS AND USAGE

Pevnar™ is indicated for active immunization of infants and toddlers against invasive disease caused by *S. pneumoniae* due to capsular serotypes included in the vaccine (4, 6B, 9V, 14, 18C, 19F, and 23F). The routine schedule is 2, 4, 6, and 12-15 months of age. For additional information on usage, see DOSAGE AND ADMINISTRATION.

This vaccine is not intended to be used for treatment of active infection.

As with any vaccine, Pevnar™ may not protect 100% of individuals receiving the vaccine.

### CONTRAINDICATIONS

Hypersensitivity to any component of the vaccine, including diphtheria toxoid, is a contraindication to use of this vaccine.

The decision to administer or delay vaccination because of a current or recent febrile illness depends largely on the severity of the symptoms and their etiology. Although a severe or even a moderate

febrile illness is sufficient reason to postpone vaccinations, minor illnesses, such as a mild upper respiratory infection with or without low-grade fever, are not generally contraindications.<sup>27,28</sup>

### **WARNINGS**

THIS VACCINE WILL NOT PROTECT AGAINST *S. PNEUMONIAE* DISEASE OTHER THAN THAT CAUSED BY THE SEVEN SEROTYPES INCLUDED IN THE VACCINE, NOR WILL IT PROTECT AGAINST OTHER MICROORGANISMS THAT CAUSE INVASIVE INFECTION SUCH AS BACTEREMIA AND MENINGITIS.

This vaccine should not be given to infants or children with thrombocytopenia or any coagulation disorder that would contraindicate intramuscular injection unless the potential benefit clearly outweighs the risk of administration. If the decision is made to administer this vaccine to children with coagulation disorders, it should be given with caution. (See DRUG INTERACTIONS).

Immunization with Prevnar™ does not substitute for routine diphtheria immunization.

Healthcare professionals should prescribe and/or administer this product with caution to patients with a possible history of latex sensitivity since the packaging contains dry natural rubber.

### **PRECAUTIONS**

Prevnar™ is for intramuscular use only. Prevnar™ SHOULD UNDER NO CIRCUMSTANCES BE ADMINISTERED INTRAVENOUSLY. The safety and immunogenicity for other routes of administration (e.g. subcutaneous) have not been evaluated.

#### **General**

CARE IS TO BE TAKEN BY THE HEALTHCARE PROFESSIONAL FOR THE SAFE AND EFFECTIVE USE OF THIS PRODUCT.

1. PRIOR TO ADMINISTRATION OF ANY DOSE OF THIS VACCINE, THE PARENT OR GUARDIAN SHOULD BE ASKED ABOUT THE PERSONAL HISTORY, FAMILY HISTORY, AND RECENT HEALTH STATUS OF THE VACCINE RECIPIENT. THE HEALTHCARE PROFESSIONAL SHOULD ASCERTAIN PREVIOUS IMMUNIZATION HISTORY, CURRENT HEALTH STATUS, AND OCCURRENCE OF ANY SYMPTOMS AND/OR SIGNS OF AN ADVERSE EVENT AFTER PREVIOUS IMMUNIZATIONS IN THE CHILD TO BE IMMUNIZED, IN ORDER TO DETERMINE THE EXISTENCE OF ANY CONTRAINDICATION TO IMMUNIZATION WITH THIS VACCINE AND TO ALLOW AN ASSESSMENT OF RISKS AND BENEFITS.
2. BEFORE THE ADMINISTRATION OF ANY BIOLOGICAL, THE HEALTHCARE PROFESSIONAL SHOULD TAKE ALL PRECAUTIONS KNOWN FOR THE PREVENTION OF ALLERGIC OR ANY OTHER ADVERSE REACTIONS. This should include a review of the patient's history regarding possible sensitivity; the ready availability of epinephrine 1:1000 and other appropriate agents used for control of immediate allergic reactions; and a knowledge of the recent literature pertaining to use of the biological concerned, including the nature of side effects and adverse reactions that may follow its use.
3. Children with impaired immune responsiveness, whether due to the use of immunosuppressive therapy (including irradiation, corticosteroids, antimetabolites, alkylating agents, and cytotoxic agents), a genetic defect, HIV infection, or other causes, may have reduced antibody response to active immunization.<sup>27,28,29</sup> (See DRUG INTERACTIONS).
4. The use of pneumococcal conjugate vaccine does not replace the use of 23-valent pneumococcal polysaccharide vaccine in children  $\geq 24$  months of age with sickle cell disease, asplenia, HIV infection, chronic illness or who are immunocompromised. Data on sequential vaccination with Prevnar™ followed by 23-valent pneumococcal polysaccharide vaccine are limited. In a randomized study, 23 children  $\geq 2$  years of age with sickle cell disease were administered either 2 doses of Prevnar™ followed by a dose of polysaccharide vaccine or a single dose of polysaccharide vaccine alone. In this small study, safety and immune responses with the combined schedule were similar to polysaccharide vaccine alone.<sup>30</sup>
5. Since this product is a suspension containing an aluminum adjuvant, shake vigorously immediately prior to use to obtain a uniform suspension prior to withdrawing the dose.

6. A separate sterile syringe and needle or a sterile disposable unit should be used for each individual to prevent transmission of hepatitis or other infectious agents from one person to another. Needles should be disposed of properly and should not be recapped.
7. Special care should be taken to prevent injection into or near a blood vessel or nerve.
8. Healthcare professionals should prescribe and/or administer this product with caution to patients with a possible history of latex sensitivity since the packaging contains dry natural rubber.

### **Information for Parents or Guardians**

Prior to administration of this vaccine, the healthcare professional should inform the parent, guardian, or other responsible adult of the potential benefits and risks to the patient (see ADVERSE REACTIONS and WARNINGS sections), and the importance of completing the immunization series unless contraindicated. Parents or guardians should be instructed to report any suspected adverse reactions to their healthcare professional. The healthcare professional should provide vaccine information statements prior to each vaccination.

### **DRUG INTERACTIONS**

Children receiving therapy with immunosuppressive agents (large amounts of corticosteroids, antimetabolites, alkylating agents, cytotoxic agents) may not respond optimally to active immunization.<sup>28,29,31,32</sup> (See PRECAUTIONS, General).

As with other intramuscular injections, Prevnar™ should be given with caution to children on anti-coagulant therapy.

### **Simultaneous Administration with Other Vaccines**

During clinical studies, Prevnar™ was administered simultaneously with DTP-HbOC or DTaP and HbOC, OPV or IPV, Hep B vaccines, MMR, and Varicella vaccine. Thus, the safety experience with Prevnar™ reflects the use of this product as part of the routine immunization schedule.<sup>19,20,22,23,25</sup>

The immune response to routine vaccines when administered with Prevnar™ (at separate sites) was assessed in 3 clinical studies in which there was a control group for comparison. Results for the concurrent immunizations in infants are shown in Table 5 and for toddlers in Table 6.

Enhancement of antibody response to HbOC in the infant series was observed. Some suppression of *Haemophilus influenzae* type b (Hib) response was seen at the 4th dose, but over 97% of children achieved titers  $\geq 1$   $\mu\text{g/mL}$ . Although some inconsistent differences in response to pertussis antigens were observed, the clinical relevance is unknown. The response to 2 doses of IPV given concomitantly with Prevnar™, assessed 3 months after the second dose, was equivalent to controls for poliovirus Types 2 and 3, but lower for Type 1. MMR and Varicella immunogenicity data from controlled clinical trials with concurrent administration of Prevnar™ are not available.

**TABLE 5**  
**Concurrent Administration of Prevnar™ With Other Vaccines to Infants in**  
**Non-Efficacy Studies<sup>20,23</sup>**

Antigen*	GMC*		% Responders†		Study	Vaccine Schedule‡ (mo.)	N	
	Prevnar™	Control§	Prevnar™	Control§			Prevnar™	Control§
Hib	6.2	4.4	99.5, 88.3	97.0, 88.1	118-12	2,4,6	214	67
Diphtheria	0.9	0.8	100	97.0				
Tetanus	3.5	4.1 <sup>  </sup>	100	100				
PT	19.1	17.8	74.0	69.7				
FHA	43.8	46.7	66.4	69.7				
Pertactin	40.1	50.9	65.6	77.3				
Fimbriae 2	3.3	4.2	44.7	62.5 <sup>  </sup>				
Hib	11.9	7.8 <sup>  </sup>	100, 96.9	98.8, 92.8	118-16	2,4,6	159	83
Hep B	--	--	99.4	96.2	118-16	0,2,6	156	80
IPV Type 1	--	--	89.0	93.6 <sup>  </sup>	118-16	2,4	156	80
Type 2	--	--	94.2	93.6				
Type 3	--	--	83.8	80.8				

\* Hib vaccine was HibTITER®, DTaP vaccine was Acel-Imune®. Hib (µg/mL); Dip, Tet (IU/mL); Pertussis Antigens (PT, FHA, Ptn, Fim) (units/mL).

† Responders = Hib (≥0.15 µg/mL, ≥1.0 µg/mL); Dip, Tet (≥0.1 IU/mL); Pertussis Antigens (PT, FHA, Ptn, Fim) [4-fold rise]; IPV (≥1:10); Hep B (≥10 mIU/mL).

‡ Schedule for concurrently administered vaccines; Prevnar™ administered at 2, 4, 6 mos.; blood for antibody assessment attained 1 month after third dose, except for IPV (3 months post-immunization).

§ Concurrent vaccines only.

|| p<0.05 when Prevnar™ compared to control group using the following tests: ANCOVA for GMCs in 118-12; ANOVA for GMCs in 118-16; and Fisher's Exact test for % Responders in 118-12.

¶ Lower bound of 90% CI of difference >10%.

**TABLE 6**  
**Concurrent Administration of Prevnar™ With Other Vaccines to Toddlers in a**  
**Non-Efficacy Study<sup>22</sup>**

Antigen*	GMC*		% Responders†		Study‡	Vaccine Schedule§ (mo.)	N	
	Prevnar™	Control	Prevnar™	Control			Prevnar™	Control
Hib	22.7	47.9¶	100, 97.9	100, 100	118-7	12-15	47	26
Diphtheria	2.0	3.2¶	100	100				
Tetanus	14.4	18.8	100	100				
PT	68.6	121.2¶	68.1	73.1				
FHA	29.0	48.2¶	68.1	84.6				
Pertactin	84.4	83.0	83.0	96.2				
Fimbriae 2	5.2	3.8	63.8	50.0				

\* Hib vaccine was HibTITER®, DTaP vaccine was Acel-Imune®. Hib (µg/mL); Dip, Tet (IU/mL); Pertussis Antigens (PT, FHA, Ptn, Fim) (units/mL).

† Responders = Hib (≥0.15 µg/mL, ≥1.0 µg/mL); Dip, Tet (≥0.1 IU/mL); Pertussis Antigens (PT, FHA, Ptn, Fim) [4-fold rise].

‡ Children received a primary series of DTP-HbOC (Tetramune®).

§ Blood for antibody assessment obtained 1 month after dose.

|| Concurrent vaccines only.

¶ p<0.05 when Prevnar™ compared to control group using a two-sample t-test.

### **CARCINOGENESIS, MUTAGENESIS, IMPAIRMENT OF FERTILITY**

Prevnar™ has not been evaluated for any carcinogenic or mutagenic potential, or impairment of fertility.

### **PREGNANCY**

#### **Pregnancy Category C**

Animal reproductive studies have not been conducted with this product. It is not known whether Prevnar™ can cause fetal harm when administered to a pregnant woman or whether it can affect reproductive capacity. This vaccine is not recommended for use in pregnant women.

#### **Nursing Mothers**

It is not known whether vaccine antigens or antibodies are excreted in human milk. This vaccine is not recommended for use in a nursing mother.

### **PEDIATRIC USE**

Prevnar™ has been shown to be usually well-tolerated and immunogenic in infants. The safety and effectiveness of Prevnar™ in children below the age of 6 weeks have not been established. Immune responses elicited by Prevnar™ among infants born prematurely have not been studied. See DOSAGE AND ADMINISTRATION for the recommended pediatric dosage.

### **GERIATRIC USE**

This vaccine is NOT recommended for use in adult populations. It is not to be used as a substitute for the pneumococcal polysaccharide vaccine, in geriatric populations.

### **ADVERSE REACTIONS**

The majority of the safety experience with Prevnar™ comes from the NCKP Efficacy Trial in which 17,066 infants received 55,352 doses of Prevnar™, along with other routine childhood vaccines through April 1998 (see CLINICAL PHARMACOLOGY section). The number of Prevnar™ recipi-

ents in the safety analysis differs from the number included in the efficacy analysis due to the different lengths of follow-up for these study endpoints. Safety was monitored in this study using several modalities. Local reactions and systemic events occurring within 48 hours of each dose of vaccine were ascertained by scripted telephone interview on a randomly selected subset of approximately 3,000 children in each vaccine group. The rate of relatively rare events requiring medical attention was evaluated across all doses in all study participants using automated databases. Specifically, rates of hospitalizations within 3, 14, 30, and 60 days of immunization, and of emergency room visits within 3, 14, and 30 days of immunization were assessed and compared between vaccine groups for each diagnosis. Seizures within 3 and 30 days of immunization were ascertained across multiple settings (hospitalizations, emergency room or clinic visits, telephone interviews). Deaths and SIDS were ascertained through April 1999. Hospitalizations due to diabetes, autoimmune disorders, and blood disorders were ascertained through August 1999. In Tables 7 and 8, the rate of local reactions at the Prevnar™ injection site is compared at each dose to the DTP or DTaP injection site in the same children.

**TABLE 7**  
**Percentage of Subjects Reporting Local Reactions Within 2 Days Following Immunization**  
**With Prevnar™ and DTP-HbOC\* Vaccines at 2, 4, 6, and 12-15 Months of Age<sup>19</sup>**

Reaction	Dose 1		Dose 2		Dose 3		Dose 4	
	<u>Prevnar™</u> Site	<u>DTP- HbOC</u> Site†	<u>Prevnar™</u> Site	<u>DTP- HbOC</u> Site†	<u>Prevnar™</u> Site	<u>DTP- HbOC</u> Site†	<u>Prevnar™</u> Site	<u>DTP- HbOC</u> Site†
	N=2890	N=2890	N=2725	N=2725	N=2538	N=2538	N=599	N=599
Erythema								
Any	12.4	21.9	14.3	25.1	15.2	26.5	12.7	23.4
> 2.4 cm	1.2	4.6	1.0	2.9	2.0	4.4	1.7	6.4
Induration								
Any	10.9	22.4	12.3	23.0	12.8	23.3	11.4	20.5
> 2.4 cm	2.6	7.2	2.4	5.6	2.9	6.7	2.8	7.2
Tenderness								
Any	28.0	36.4	25.2	30.5	25.6	32.8	36.5	45.1
Interfered with limb movement	7.9	10.7	7.4	8.4	7.8	10.0	18.5	22.2

\* If Hep B vaccine was administered simultaneously, it was administered into the same limb as the DTP-HbOC vaccine. If reactions occurred at either or both sites on that limb, the more severe reaction was recorded.

† p<0.05 when Prevnar™ site compared to the DTP-HbOC site using the sign test.

**TABLE 8**  
**Percentage of Subjects Reporting Local Reactions Within 2 Days Following Immunization**  
**With Prevnar™\* and DTaP Vaccines† at 2, 4, 6, and 12-15 Months of Age<sup>19</sup>**

Reaction	Dose 1		Dose 2		Dose 3		Dose 4	
	Prevnar™ Site	DTaP Site	Prevnar™ Site	DTaP Site	Prevnar™ Site	DTaP Site	Prevnar™ Site	DTaP Site‡
	N=693	N=693	N=526	N=526	N=422	N=422	N=165	N=165
Erythema								
Any	10.0	6.7§	11.6	10.5	13.8	11.4	10.9	3.6§
> 2.4 cm	1.3	0.4§	0.6	0.6	1.4	1.0	3.6	0.6
Induration								
Any	9.8	6.6§	12.0	10.5	10.4	10.4	12.1	5.5§
> 2.4 cm	1.6	0.9	1.3	1.7	2.4	1.9	5.5	1.8
Tenderness								
Any	17.9	16.0	19.4	17.3	14.7	13.1	23.3	18.4
Interfered with limb movement	3.1	1.8§	4.1	3.3	2.9	1.9	9.2	8.0

\* HbOC was administered in the same limb as Prevnar™. If reactions occurred at either or both sites on that limb, the more severe reaction was recorded.

† If Hep B vaccine was administered simultaneously, it was administered into the same limb as DTaP. If reactions occurred at either or both sites on that limb, the more severe reaction was recorded.

‡ Subjects may have received DTP or a mixed DTP/DTaP regimen for the primary series. Thus, this is the 4th dose of a pertussis vaccine, but not a 4th dose of DTaP.

§ p<0.05 when Prevnar™ site compared to DTaP site using the sign test.

Table 9 presents the rates of local reactions in previously unvaccinated older infants and children.

**TABLE 9**  
**Percentage of Subjects Reporting Local Reactions Within 3 Days of Immunization in Infants and Children from 7 Months Through 9 Years of Age<sup>26</sup>**

Age at 1st Vaccination	7 - 11 Mos.						12 - 23 Mos.			24 - 35 Mos.	36 - 59 Mos.	5 - 9 Yrs.
	118-12			118-16			118-9*	118-18		118-18	118-18	118-18
Dose Number	1	2	3†	1	2	3†	1	1	2	1	1	1
Number of Subjects	54	51	24	81	76	50	60	114	117	46	48	49
Reaction												
Erythema												
Any	16.7	11.8	20.8	7.4	7.9	14.0	48.3	10.5	9.4	6.5	29.2	24.2
> 2.4 cm‡	1.9	0.0	0.0	0.0	0.0	0.0	6.7	1.8	1.7	0.0	8.3	7.1
Induration												
Any	16.7	11.8	8.3	7.4	3.9	10.0	48.3	8.8	6.0	10.9	22.9	25.5
> 2.4 cm‡	3.7	0.0	0.0	0.0	0.0	0.0	3.3	0.9	0.9	2.2	6.3	9.3
Tenderness												
Any	13.0	11.8	12.5	8.6	10.5	12.0	46.7	25.7	26.5	41.3	58.3	82.8
Interfered with limb movement§	1.9	2.0	4.2	1.2	1.3	0.0	3.3	6.2	8.5	13.0	20.8	39.4

\* For 118-9, 2 of 60 subjects were ≥24 months of age.

† For 118-12, dose 3 was administered at 15 - 18 mos. of age. For 118-16, dose 3 was administered at 12 - 15 mos. of age.

‡ For 118-16 and 118-18, ≥2 cm.

§ Tenderness interfering with limb movement.

Tables 10 and 11 present the rates of systemic events observed in the efficacy study when Prevnar™ was administered concomitantly with DTP or DTaP.

**TABLE 10**  
**Percentage of Subjects\* Reporting Systemic Events Within 2 Days Following Immunization With Prevnar™ or Control† Vaccine Concurrently With DTP-HbOC Vaccine at 2, 4, 6, and 12-15 Months of Age<sup>18</sup>**

Reaction	Dose 1		Dose 2		Dose 3		Dose 4	
	Prevnar™	Control†	Prevnar™	Control†	Prevnar™	Control†	Prevnar™	Control†
	N=2998	N=2982	N=2788	N=2761	N=2596	N=2591	N=709	N=733
Fever								
≥ 38.0°C	33.4	28.7‡	34.7	27.4‡	40.6	32.4‡	41.9	36.9
> 39.0°C	1.3	1.3	3.0	1.6‡	5.3	3.4‡	4.5	4.5
Irritability	71.3	67.9‡	69.4	63.8‡	68.9	61.6‡	72.8	65.8‡
Drowsiness	49.2	50.6	32.5	33.6	25.9	23.4‡	21.3	22.7
Restless Sleep	18.1	17.9	27.3	24.3‡	33.3	30.1‡	29.9	28.0
Decreased Appetite	24.7	23.6	22.8	20.3‡	27.7	25.6	33.0	27.4‡
Vomiting	17.9	14.9‡	16.2	14.4	15.5	12.7‡	9.6	6.8
Diarrhea	12.0	10.7	10.9	9.9	11.5	10.4	12.1	11.2
Rash or Hives	0.7	0.6	0.8	0.8	1.4	1.1	1.4	0.8

\* Approximately 90% of subjects received prophylactic or therapeutic antipyretics within 48 hours of each dose.

† Investigational meningococcal group C conjugate vaccine (MnCC).

‡ p<0.05 when Prevnar™ compared to control group using a Chi-Square test.

**TABLE 11**  
**Percentage of Subjects\* Reporting Systemic Events Within 2 Days Following**  
**Immunization With Prevnar™ or Control† Vaccine Concurrently With DTaP**  
**Vaccine at 2, 4, 6, and 12-15 Months of Age<sup>18</sup>**

Reaction	Dose 1		Dose 2		Dose 3		Dose 4	
	Prevnar™	Control†	Prevnar™	Control†	Prevnar™	Control†	Prevnar™	Control†
	N=710	N=711	N=559	N=508	N=461	N=414	N=224	N=230
Fever								
≥ 38.0°C	15.1	9.4§	23.9	10.8§	19.1	11.8§	21.0	17.0
> 39.0°C	0.9	0.3	2.5	0.8§	1.7	0.7	1.3	1.7
Irritability	48.0	48.2	58.7	45.3§	51.2	44.8	44.2	42.6
Drowsiness	40.7	42.0	25.6	22.8	19.5	21.9	17.0	16.5
Restless Sleep	15.3	15.1	20.2	19.3	25.2	19.0§	20.2	19.1
Decreased Appetite	17.0	13.5	17.4	13.4	20.7	13.8§	20.5	23.1
Vomiting	14.6	14.5	16.8	14.4	10.4	11.6	4.9	4.8
Diarrhea	11.9	8.4§	10.2	9.3	8.3	9.4	11.6	9.2
Rash or Hives	1.4	0.3§	1.3	1.4	0.4	0.5	0.5	1.7

\* Approximately 75% of subjects received prophylactic or therapeutic antipyretics within 48 hours of each dose.

† Investigational meningococcal group C conjugate vaccine (MnCC).

‡ Most of these children had received DTP for the primary series. Thus, this is a 4th dose of a pertussis vaccine, but not of DTaP.

§ p<0.05 when Prevnar™ compared to control group using a Chi-Square test.

Table 12 presents results from a second study (Manufacturing Bridging Study) conducted at Northern California and Denver Kaiser sites, in which children were randomized to receive one of three lots of Prevnar™ with concomitant vaccines including DTaP, or the same concomitant vaccines alone. Information was ascertained by scripted telephone interview, as described above.

**TABLE 12**  
**Percentage of Subjects\* Reporting Systemic Reactions Within 3 Days Following Immunization With Prevnar™, DTaP, HbOC, Hep B, and IPV vs. Control† In Manufacturing Bridging Study<sup>20</sup>**

Reaction	Dose 1		Dose 2		Dose 3	
	Prevnar™	Control†	Prevnar™	Control†	Prevnar™	Control†
	N=498	N=108	N=452	N=99	N=445	N=89
Fever						
≥ 38.0°C	21.9	10.2‡	33.6	17.2‡	28.1	23.6
> 39.0°C	0.8	0.9	3.8	0.0	2.2	0.0
Irritability	59.7	60.2	65.3	52.5‡	54.2	50.6
Drowsiness	50.8	38.9‡	30.3	31.3	21.2	20.2
Decreased Appetite	19.1	15.7	20.6	11.1‡	20.4	9.0‡

\* Approximately 72% of subjects received prophylactic or therapeutic antipyretics within 48 hours of each dose.

† Control group received concomitant vaccines only in the same schedule as the Prevnar™ group (DTaP, HbOC at dose 1, 2, 3; IPV at doses 1 and 2; Hep B at doses 1 and 3).

‡ p<0.05 when Prevnar™ compared to control group using Fisher's Exact test.

Fever (≥ 38.0°C) within 48 hours of a vaccine dose was reported by a greater proportion of subjects who received Prevnar™, compared to control (investigational meningococcal group C conjugate vaccine [MnCC]), after each dose when administered concurrently with DTP-HbOC or DTaP in the efficacy study. In the Manufacturing Bridging Study, fever within 48-72 hours was also reported more commonly after each dose compared to infants in the control group who received only recommended vaccines. When administered concurrently with DTaP in either study, fever rates among Prevnar™ recipients ranged from 15% to 34%, and were greatest after the 2nd dose.

Table 13 presents the frequencies of systemic reactions in previously unvaccinated older infants and children.

**TABLE 13**  
**Percentage of Subjects Reporting Systemic Reactions Within 3 Days of Immunization**  
**in Infants and Children from 7 Months Through 9 Years of Age<sup>26</sup>**

Age at 1st Vaccination	7 - 11 Mos.						12 - 23 Mos.			24 - 35 Mos.	36 - 59 Mos.	5 - 9 Yrs.
	118-12			118-16			118-9*	118-18		118-18	118-18	118-18
Dose Number	1	2	3†	1	2	3†	1	1	2	1	1	1
Number of Subjects	54	51	24	85	80	50	60	120	117	47	52	100
<b>Reaction</b>												
Fever												
≥ 38.0°C	20.8	21.6	25.0	17.6	18.8	22.0	36.7	11.7	6.8	14.9	11.5	7.0
> 39.0°C	1.9	5.9	0.0	1.6	3.9	2.6	0.0	4.4	0.0	4.2	2.3	1.2
Fussiness	29.6	39.2	16.7	54.1	41.3	38.0	40.0	37.5	36.8	46.8	34.6	29.3
Drowsiness	11.1	17.6	16.7	24.7	16.3	14.0	13.3	18.3	11.1	12.8	17.3	11.0
Decreased Appetite	9.3	15.7	0.0	15.3	15.0	30.0	25.0	20.8	16.2	23.4	11.5	9.0

\* For 118-9, 2 of 60 subjects were ≥24 months of age.

† For 118-12, dose 3 was administered at 15 - 18 mos. of age. For 118-16, dose 3 was administered at 12 - 15 mos. of age.

Of the 17,066 subjects who received at least one dose of Prevnar™ in the efficacy trial, there were 24 hospitalizations (for 29 diagnoses) within 3 days of a dose from October 1995 through April 1998. Diagnoses were as follows: bronchiolitis (5); congenital anomaly (4); elective procedure, UTI (3 each); acute gastroenteritis, asthma, pneumonia (2 each); aspiration, breath holding, influenza, inguinal hernia repair, otitis media, febrile seizure, viral syndrome, well child/reassurance (1 each). There were 162 visits to the emergency room (for 182 diagnoses) within 3 days of a dose from October 1995 through April 1998. Diagnoses were as follows: febrile illness (20); acute gastroenteritis (19); trauma, URI (16 each); otitis media (15); well child (13); irritable child, viral syndrome (10 each); rash (8); croup, pneumonia (6 each); poisoning/ingestion (5); asthma, bronchiolitis (4 each); febrile seizure, UTI (3 each); thrush, wheezing, breath holding, choking, conjunctivitis, inguinal hernia repair, pharyngitis (2 each); colic, colitis, congestive heart failure, elective procedure, hives, influenza, ingrown toenail, local swelling, roseola, sepsis (1 each).<sup>19</sup>

One case of a hypotonic-hyporesponsive episode (HHE) was reported in the efficacy study following Prevnar™ and concurrent DTP vaccines in the study period from October 1995 through April 1998. Two additional cases of HHE were reported in four other studies and these also occurred in children who received Prevnar™ concurrently with DTP vaccine.<sup>22,25</sup>

In the Kaiser efficacy study in which 17,066 children received a total of 55,352 doses of Prevnar™ and 17,080 children received a total of 55,387 doses of the control vaccine (investiga-

tional meningococcal group C conjugate vaccine [MnCC]), seizures were reported in 8 Prevnar™ recipients and 4 control vaccine recipients within 3 days of immunization from October 1995 through April 1998. Of the 8 Prevnar™ recipients, 7 received concomitant DTP-containing vaccines and one received DTaP. Of the 4 control vaccine recipients, 3 received concomitant DTP-containing vaccines and one received DTaP.<sup>19</sup> In the other 4 studies combined, in which 1,102 children were immunized with 3,347 doses of Prevnar™ and 408 children were immunized with 1,310 doses of control vaccine (either investigational meningococcal group C conjugate vaccine [MnCC] or concurrent vaccines), there was one seizure event reported within 3 days of immunization.<sup>23</sup> This subject received Prevnar™ concurrent with DTaP vaccine.

Twelve deaths (5 SIDS and 7 with clear alternative cause) occurred among subjects receiving Prevnar™, of which 11 (4 SIDS and 7 with clear alternative cause) occurred in the Kaiser efficacy study from October 1995 until April 20, 1999. In comparison, 21 deaths (8 SIDS, 12 with clear alternative cause and one SIDS-like death in an older child), occurred in the control vaccine group during the same time period in the efficacy study.<sup>19,20</sup> The number of SIDS deaths in the efficacy study from October 1995 until April 20, 1999 was similar to or lower than the age and season-adjusted expected rate from the California State data from 1995-1997 and are presented in Table 14.

**TABLE 14**  
**Age and Season-Adjusted Comparison of SIDS Rates in the NCKP Efficacy Trial**  
**With the Expected Rate from the California State Data for 1995-1997<sup>19</sup>**

Vaccine	< One Week After Immunization		≤ Two Weeks After Immunization		≤ One Month After Immunization		≤ One Year After Immunization	
	Exp	Obs	Exp	Obs	Exp	Obs	Exp	Obs
Prevnar™	1.06	1	2.09	2	4.28	2	8.08	4
Control*	1.06	2	2.09	3†	4.28	3†	8.08	8†

\* Investigational meningococcal group C conjugate vaccine (MnCC).

† Does not include one additional case of SIDS-like death in a child older than the usual SIDS age (448 days).

In a review of all hospitalizations that occurred between October 1995 and August 1999 in the efficacy study for the specific diagnoses of aplastic anemia, autoimmune disease, autoimmune hemolytic anemia, diabetes mellitus, neutropenia, and thrombocytopenia, the numbers of such cases were either equal to or less than the expected numbers based on the 1995 Kaiser Vaccine Safety Data Link (VSD) data set.

Overall, the safety of Prevnar™ was evaluated in a total of five clinical studies in which 18,168 infants and children received a total of 58,699 doses of vaccine at 2, 4, 6, and 12-15 months of age. In addition, the safety of Prevnar™ was evaluated in 560 children from 4 ancillary studies who started immunization at 7 months to 9 years of age. Tables 15 and 16 summarize systemic reactogenicity data within 2 or 3 days across 4,748 subjects (13,039 infant doses and 1,706 toddler doses) for whom these data were collected and according to the pertussis vaccine administered concurrently.

**TABLE 15**  
**Overall Percentage of Doses Associated With Systemic Events Within 2 or 3 Days**  
**For Efficacy Study and All Ancillary Studies When Prevnar™ Administered To**  
**Infants As a Primary Series at 2, 4, and 6 Months of Age<sup>19,20,22,23,25</sup>**

<b>Systemic Event</b>	<b>Prevnar™ Concurrently With DTP-HbOC (9,191 Doses) *</b>	<b>Prevnar™ Concurrently With DTaP and HbOC (3,848 Doses) †</b>	<b>DTaP and HbOC Control (538 Doses) ‡</b>
<b>Fever</b>			
≥ 38.0°C	35.6	21.1	<b>14.2</b>
> 39.0°C	3.1	1.8	<b>0.4</b>
<b>Irritability</b>	69.1	52.5	<b>45.2</b>
<b>Drowsiness</b>	36.9	32.9	<b>27.7</b>
<b>Restless Sleep</b>	25.8	20.6	<b>22.3</b>
<b>Decreased Appetite</b>	24.7	18.1	<b>13.6</b>
<b>Vomiting</b>	16.2	13.4	<b>9.8</b>
<b>Diarrhea</b>	11.4	9.8	<b>4.4</b>
<b>Rash or Hives</b>	0.9	0.6	<b>0.3</b>

\* Total from which reaction data are available varies between reactions from 8,874-9,191 doses. Data from studies 118-3, 118-7, 118-8.

† Total from which reaction data are available varies between reactions from 3,121-3,848 doses. Data from studies 118-8, 118-12, 118-16.

‡ Total from which reaction data are available varies between reactions from 295-538 doses. Data from studies 118-12 and 118-16.

**TABLE 16**  
**Overall Percentage of Doses Associated With Systemic Events Within 2 or 3 Days**  
**For Efficacy Study and All Ancillary Studies When Prevnar™ Administered To**  
**Toddlers as a Fourth Dose At 12 to 15 Months of Age<sup>19,22</sup>**

<b>Systemic Event</b>	<b>Prevnar™ Concurrently With DTP-HbOC (709 Doses)*</b>	<b>Prevnar™ Concurrently With DTaP and HbOC (270 Doses)†</b>	<b>Prevnar™ Only No Concurrent Vaccines (727 Doses)‡</b>
<b>Fever</b>			
<b>≥ 38.0°C</b>	41.9	19.6	<b>13.4</b>
<b>&gt; 39.0°C</b>	4.5	1.5	<b>1.2</b>
<b>Irritability</b>	72.8	45.9	<b>45.8</b>
<b>Drowsiness</b>	21.3	17.5	<b>15.9</b>
<b>Restless Sleep</b>	29.9	21.2	<b>21.2</b>
<b>Decreased Appetite</b>	33.0	21.1	<b>18.3</b>
<b>Vomiting</b>	9.6	5.6	<b>6.3</b>
<b>Diarrhea</b>	12.1	13.7	<b>12.8</b>
<b>Rash or Hives</b>	1.4	0.7	<b>1.2</b>

\* Total from which reaction data are available varies between reactions from 706-709 doses. Data from study 118-8.

† Total from which reaction data are available varies between reactions from 269-270 doses. Data from studies 118-7 and 118-8.

‡ Total from which reaction data are available varies between reactions from 725-727 doses. Data from studies 118-7 and 118-8.

With vaccines in general, including Prevnar™, it is not uncommon for patients to note within 48 to 72 hours at or around the injection site the following minor reactions: edema; pain or tenderness; redness, inflammation or skin discoloration; mass; or local hypersensitivity reaction. Such local reactions are usually self-limited and require no therapy.

As with other aluminum-containing vaccines, a nodule may occasionally be palpable at the injection site for several weeks.<sup>33</sup>

### **ADVERSE EVENT REPORTING**

Any suspected adverse events following immunization should be reported by the healthcare professional to the US Department of Health and Human Services (DHHS). The National Vaccine Injury Compensation Program requires that the manufacturer and lot number of the vaccine administered be recorded by the healthcare professional in the vaccine recipient's permanent medical record (or in a permanent office log or file), along with the date of administration of the vaccine and the name, address, and title of the person administering the vaccine.

The US DHHS has established the Vaccine Adverse Event Reporting System (VAERS) to accept all

reports of suspected adverse events after the administration of any vaccine including, but not limited to, the reporting of events required by the National Childhood Vaccine Injury Act of 1986. The FDA web site is: <http://www.fda.gov/cber/vaers/vaers.htm>.

The VAERS toll-free number for VAERS forms and information is 800-822-7967.<sup>34</sup>

### **DOSAGE AND ADMINISTRATION**

**For intramuscular injection only. Do not inject intravenously.**

The dose is 0.5 mL to be given intramuscularly.

**Since this product is a suspension containing an adjuvant, shake vigorously immediately prior to use to obtain a uniform suspension in the vaccine container. The vaccine should not be used if it cannot be resuspended.**

After shaking, the vaccine is a homogeneous, white suspension.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration (see DESCRIPTION). This product should not be used if particulate matter or discoloration is found.

The vaccine should be injected intramuscularly. The preferred sites are the anterolateral aspect of the thigh in infants or the deltoid muscle of the upper arm in toddlers and young children. The vaccine should not be injected in the gluteal area or areas where there may be a major nerve trunk and/or blood vessel. Before injection, the skin at the injection site should be cleansed and prepared with a suitable germicide. After insertion of the needle, aspirate and wait to see if any blood appears in the syringe, which will help avoid inadvertent injection into a blood vessel. If blood appears, withdraw the needle and prepare for a new injection at another site.

### **Vaccine Schedule**

For infants, the immunization series of Prevnar™ consists of three doses of 0.5 mL each, at approximately 2-month intervals, followed by a fourth dose of 0.5 mL at 12-15 months of age. The customary age for the first dose is 2 months of age, but it can be given as young as 6 weeks of age. The recommended dosing interval is 4 to 8 weeks. The fourth dose should be administered at least 2 months after the third dose.

### **Previously Unvaccinated Older Infants and Children**

For previously unvaccinated older infants and children, who are beyond the age of the routine infant schedule, the following schedule applies:<sup>26</sup>

<b>Age at First Dose</b>	<b>Total Number of 0.5 mL Doses</b>
7-11 months of age	3*
12-23 months of age	2†
≥ 24 months through 9 years of age	1

\* 2 doses at least 4 weeks apart; third dose after the one-year birthday, separated from the second dose by at least 2 months.

† 2 doses at least 2 months apart.

(See CLINICAL PHARMACOLOGY section for the limited available immunogenicity data and ADVERSE EVENTS section for limited safety data corresponding to the previously noted vaccination schedule for older children).

Safety and immunogenicity data are either limited or not available for children in specific high risk groups for invasive pneumococcal disease (e.g. persons with sickle cell disease, asplenia, HIV-infected).

### **HOW SUPPLIED**

Vial, 1 Dose (5 per package) - NDC 0005-1970-67

CPT Code 90669

## STORAGE

DO NOT FREEZE. STORE REFRIGERATED, AWAY FROM FREEZER COMPARTMENT, AT 2°C TO 8°C (36°F TO 46°F).

## REFERENCES

1. Schuchat A, Robinson K, Wenger JD, et al. Bacterial meningitis in the United States in 1995. *N Engl J Med*. 1997; 337:970-6.
2. Zangwill KM, Vadheim CM, Vannier AM, et al. Epidemiology of invasive pneumococcal disease in Southern California: implications for the design and conduct of a pneumococcal conjugate vaccine efficacy trial. *J Infect Dis*. 1996; 174:752-9.
3. Pastor P, Medley F, Murphy T. Invasive pneumococcal disease in Dallas County, Texas: results from population-based surveillance in 1995. *Clin Infect Dis*. 1998; 26:590-5.
4. Hofmann J, Cetron MS, Farley MM, et al. The prevalence of drug-resistant *Streptococcus pneumoniae* in Atlanta. *N Engl J Med*. 1995; 333:481-515.
5. Breiman R, Spika J, Navarro V, et al. Pneumococcal bacteremia in Charleston County, South Carolina. *Arch Intern Med*. 1990; 150:1401-5.
6. Plouffe J, Breiman R, Facklam R. Franklin County Study Group. Bacteremia with *Streptococcus pneumoniae* in adults-implications for therapy and prevention. *JAMA*. 1996; 275:194-8.
7. Levine O, Farley M, Harrison LH, et al. Risk factors for invasive pneumococcal disease in children: a population-based case-control study in North America. *Pediatrics*. 1999; 103:1-5.
8. Kaplan SL, Mason EO, Barson WJ, et al. Three-year multicenter surveillance of systemic pneumococcal infections in children. *Pediatrics*. 1998; 102:538-44.
9. Arditi M, Edward M, Bradley J, et al. Three-year multicenter surveillance of pneumococcal meningitis in children: clinical characteristics and outcome related to penicillin susceptibility and dexamethasone use. *Pediatrics*. 1998; 102:1087-97.
10. Bluestone CD, Stephenson BS, Martin LM. Ten-year review of otitis media pathogens. *Pediatr Infect Dis J*. 1992; 11:S7-S11.
11. Giebink GS. The microbiology of otitis media. *Pediatr Infect Dis J*. 1989; 8:S18-S20.
12. Butler JC, Breiman RF, Lipman HB, et al. Serotype distribution of *Streptococcus pneumoniae* infections among preschool children in the United States, 1978-1994: implications for development of a conjugate vaccine. *J Infect Dis*. 1995; 171:885-9.
13. Paisley JW, Lauer BA, McIntosh K, et al. Pathogens associated with acute lower respiratory tract infection in young children. *Pediatr Infect Dis J*. 1984; 3:14-9.
14. Heiskanen-Kosma T, Korppi M, Jokinen C, et al. Etiology of childhood pneumonia: serologic results of a prospective, population-based study. *Pediatr Infect Dis J*. 1998; 17:986-91.
15. American Academy of Pediatrics Committee on Infectious Diseases. Therapy for children with invasive pneumococcal infections. *Pediatrics*. 1997; 99:289-300.
16. Sniadack DH, Schwart B, Lipman H, et al. Potential interventions for the prevention of childhood pneumonia: geographic and temporal differences in serotype and serogroup distribution of sterile site pneumococcal isolates from children-implications for vaccine strategies. *Pediatr Infect Dis J*. 1995; 14:503-10.
17. Butler JC, Hoffman J, Cetron MS, et al. The continued emergence of drug-resistant *Streptococcus pneumoniae* in the United States. An Update from the Centers for Disease Control and Prevention's Pneumococcal Sentinel Surveillance System. *J Infect Dis*. 1996; 174:986-93.
18. Black S, Shinefield H, Ray P, et al. Efficacy of Heptavalent Conjugate Pneumococcal Vaccine (Lederle Laboratories) in 37,000 Infants and Children. Results of the Northern California Kaiser Permanente Efficacy Trial. 38th ICAAC, San Diego, California, September 24-27, 1998.
19. Lederle Laboratories, Data on File: D118-P8.

20. Lederle Laboratories, Data on File: D118-P16.
21. Lederle Laboratories, Data on File: D118-P8 Addendum DTaP Immunogenicity.
22. Shinefield HR, Black S, Ray P. Safety and immunogenicity of heptavalent pneumococcal CRM<sub>197</sub> conjugate vaccine in infants and toddlers. *Pediatr Infect Dis J*. 1999; 18:757-63.
23. Lederle Laboratories, Data on File: D118-P12.
24. Rennels MD, Edwards KM, Keyserling HL, et al. Safety and immunogenicity of heptavalent pneumococcal vaccine conjugated to CRM<sub>197</sub> in United States infants. *Pediatrics*. 1998; 101(4):604-11.
25. Lederle Laboratories, Data on File: D118-P3.
26. Lederle Laboratories, Data on File: Integrated Summary on Catch-Up.
27. Report of the Committee on Infectious Diseases 24th Edition. Elk Grove Village, IL: American Academy of Pediatrics. 1997; 31-3.
28. Update: Vaccine Side Effects, Adverse Reactions, Contraindications, and Precautions. *MMWR*. 1996; 45(RR-12):1-35.
29. Recommendations of the Advisory Committee on Immunization Practices (ACIP): use of vaccines and immunoglobulins in persons with altered immunocompetence. *MMWR*. 1993; 43(RR-4):1-18.
30. Vernacchio L, Neufeld EJ, MacDonald K, et al. Combined schedule of 7-valent pneumococcal conjugate vaccine followed by 23-valent pneumococcal vaccine in children and young adults with sickle cell disease. *J Pediatr*. 1998; 103:275-8.
31. Immunization of children infected with human immunodeficiency virus-supplementary ACIP statement. *MMWR*. 1988; 37(12):181-83.
32. Center for Disease Control. General recommendations on immunization. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*. 1994; 43(RR01):1-38.
33. Fawcett HA, Smith NP. Injection-site granuloma due to aluminum. *Archives Dermatology*. 1984; 120:1318-22.
34. Vaccines Adverse Event Reporting System – United States. *MMWR*. 1990; 39:730-3.

Manufactured by:

**LEDERLE LABORATORIES**

Division American Cyanamid Company

Pearl River, NY 10965 USA

US GOVT. LICENSE NO. 17

Marketed by:

**WYETH LEDERLE**

VACCINES

Philadelphia, PA 19101

CI 6044-1

PRINTED IN USA  
Issued February 16, 2000