

Behavioral Counseling in Primary Care to Promote a Healthy Diet

Recommendations and Rationale

U.S. Preventive Services Task Force

This statement summarizes the U.S. Preventive Services Task Force (USPSTF) recommendations on counseling to promote a healthy diet in primary care patients and the supporting evidence, and it updates the 1996 recommendations contained in the *Guide to Clinical Preventive Services*, second edition.¹ Explanations of the ratings and of the strength of overall evidence are given in Appendix A and Appendix B, respectively. The complete information on which this statement is based, including evidence tables and references, is available in the Systematic Evidence Review² on this topic, which can be obtained through the USPSTF web site (www.preventiveservices.ahrq.gov) and through the National Guideline Clearinghouse™ (<http://www.guideline.gov>). The summary of the evidence and this recommendation statement are also available in print through the AHRQ Publications Clearinghouse (call 1-800-358-9295 or e-mail ahrqpubs@ahrq.gov).

This was first published in *Am J Prev Med*. 2003; 24(1):93-100.

Summary of Recommendations

The U.S. Preventive Services Task Force (USPSTF) concludes that the evidence is insufficient to recommend for or against routine behavioral counseling to promote a healthy diet in unselected patients in primary care settings.

I recommendation.

The USPSTF found fair evidence that brief, low- to medium-intensity behavioral dietary counseling in the primary care setting can produce small to medium changes in average daily intake of core components of

an overall healthy diet (especially saturated fat and fruit and vegetables) in unselected patients (see “Scientific Evidence” for discussion of patient populations and intensity of interventions). The strength of this evidence, however, is limited by reliance on self-reported diet outcomes, limited use of measures corroborating reported changes in diet, limited follow-up data beyond 6 to 12 months, and enrollment of study participants who may not be fully representative of primary care patients. In addition, there is limited evidence to assess possible harms (see “Clinical Considerations”). As a result, the USPSTF concluded that there is insufficient evidence to determine the significance and magnitude of the benefit of routine counseling to promote a healthy diet in adults. Although community-based studies have evaluated measures to reduce dietary fat intake in children, no controlled trials of routine behavioral dietary counseling for children or adolescents in the primary care setting were identified.

The USPSTF recommends intensive behavioral dietary counseling for adult patients with hyperlipidemia and other known risk factors for cardiovascular and diet-related chronic disease. Intensive counseling can be delivered by primary care clinicians or by referral to other specialists, such as nutritionists or dietitians.

B recommendation.

The USPSTF found good evidence that medium- to high-intensity counseling interventions can produce medium to large changes in average daily intake of

Corresponding Author: Alfred O. Berg, MD, MPH, Chair, U.S. Preventive Services Task Force, c/o David Atkins, MD, MPH, Chief Medical Officer, Center for Practice and Technology Assessment, Agency for Healthcare Research and Quality, 6010 Executive Boulevard, Suite 300, Rockville, MD 20852. (301) 594-4016, fax (301) 594-4027, E-mail: uspstf@ahrq.gov.

core components of a healthy diet (including saturated fat, fiber, fruit, and vegetables) among adult patients at increased risk for diet-related chronic disease. Intensive counseling interventions that have been examined in controlled trials among at-risk adult patients have combined nutrition education with behavioral dietary counseling provided by a nutritionist, dietitian, or specially trained primary care clinician (eg, physician, nurse, or nurse practitioner). The USPSTF concluded that such counseling is likely to improve important health outcomes and that benefits outweigh potential harms. No controlled trials of intensive counseling in children or adolescents that measured diet were identified.^{3,4}

Clinical Considerations

- Several brief dietary assessment questionnaires have been validated for use in the primary care setting.^{5,6} These instruments can identify dietary counseling needs, guide interventions, and monitor changes in patients' dietary patterns. However, these instruments are susceptible to the bias of the respondent. Therefore, when used to evaluate the efficacy of counseling, efforts to verify self-reported information are recommended since patients receiving dietary interventions may be more likely to report positive changes in dietary behavior than control patients.⁷⁻¹⁰
- Effective interventions combine nutrition education with behaviorally-oriented counseling to help patients acquire the skills, motivation, and support needed to alter their daily eating patterns and food preparation practices. Examples of behaviorally-oriented counseling interventions include teaching self-monitoring, training to overcome common barriers to selecting a healthy diet, helping patients to set their own goals, providing guidance in shopping and food preparation, role playing, and arranging for intra-treatment social support. In general, these interventions can be described with reference to the 5-A behavioral counseling framework¹¹: *Assess* dietary practices and related risk factors, *Advise* to change dietary practices, *Agree* on individual diet change goals, *Assist* to change dietary practices or address motivational barriers, and *Arrange* regular follow-up and support or refer to more intensive behavioral nutritional counseling (eg, medical nutrition therapy) if needed.
- Two approaches appear promising for the general population of adult patients in primary care settings: (1) medium-intensity face-to-face dietary counseling (2 to 3 group or individual sessions) delivered by a dietitian or nutritionist or by a specially trained primary care physician or nurse practitioner, and (2) lower-intensity interventions that involve 5 minutes or less of primary care provider counseling supplemented by patient self-help materials, telephone counseling, or other interactive health communications. However, more research is needed to assess the long-term efficacy of these treatments and the balance of benefits and harms.
- The largest effect of dietary counseling in asymptomatic adults has been observed with more intensive interventions (multiple sessions lasting 30 minutes or longer) among patients with hyperlipidemia or hypertension, and among others at increased risk for diet-related chronic disease. Effective interventions include individual or group counseling delivered by nutritionists, dietitians, or specially trained primary care practitioners or health educators in the primary care setting or in other clinical settings by referral. Most studies of these interventions have enrolled selected patients, many of whom had known diet-related risk factors such as hyperlipidemia or hypertension. Similar approaches may be effective with unselected adult patients, but adherence to dietary advice may be lower, and health benefits smaller, than in patients who have been told they are at higher risk for diet-related chronic disease.¹²
- Office-level systems supports (prompts, reminders, and counseling algorithms) have been found to significantly improve the delivery of appropriate dietary counseling by primary care clinicians.¹³⁻¹⁵
- Possible harms of dietary counseling have not been well defined or measured. Some have raised concerns that if patients focus only on reducing total fat intake without attention to reducing

caloric intake, an increase in carbohydrate intake (eg, reduced-fat or low-fat food products) may lead to weight gain, elevated triglyceride levels, or insulin resistance. Nationally, obesity rates have increased despite declining fat consumption, but studies did not consistently examine effects of counseling on outcomes such as caloric intake and weight.

- Little is known about effective dietary counseling for children or adolescents in the primary care setting. Most studies of nutritional interventions for children and adolescents have focused on non-clinical settings (such as schools) or have used physiologic outcomes such as cholesterol or weight rather than more comprehensive measures of a healthy diet.^{3,4}

Scientific Evidence

Epidemiology and Clinical Consequences

Consuming a healthy diet is associated with lower risks for chronic disease morbidity and mortality. Four of the 10 leading causes of death—coronary heart disease, some types of cancer, stroke, and type 2 diabetes—are associated with unhealthy diets.² The relationships between dietary patterns and health outcomes have been examined in a wide range of observational studies and randomized trials with patients at risk for diet-related chronic disease. The majority of studies show that people consuming diets that are low in fat, saturated fat, trans-fatty acids, and cholesterol and high in fruits, vegetables, and whole grain products containing fiber have lower rates of morbidity and mortality from coronary heart disease, and possibly several forms of cancer. In addition, one needs to balance calories with physical activity to maintain a healthy weight. The Dietary Guidelines for Americans¹⁶ recommend 3 to 5 daily servings of vegetables and vegetable juices, 2 to 4 daily servings of fruits and fruit juices, and 6 to 11 daily servings of grain products, depending on caloric needs. In addition, they recommend a diet that contains less than 10% of calories from saturated fat, no more than 30% of calories from total fat, and limited consumption of trans-fatty acids.

Despite well-established benefits of consuming a healthy diet, more than 80% of Americans of all ages eat fewer than the recommended number of daily servings of fruit, vegetables, and grain products and more than the recommended proportions of daily calories from saturated fat and total fat.¹⁷ In 1994-1996, 28% of people aged 2 years and older consumed at least 2 daily servings of fruit, 49% consumed at least 3 daily servings of vegetables, 51% consumed at least 6 daily servings of grain products, 36% consumed less than 10% of daily calories from saturated fat, and 33% consumed 30% or less of daily calories from total fat.¹⁷

Dietary counseling practices of primary care clinicians indicate limited attention to diet modification. In a 1999-2000 survey of U.S. adults, 33% of respondents reported past-year physician advice to eat more fruits and vegetables, and 29% reported similar advice to reduce dietary fat.¹⁸ In another recent survey, 25% of adult patients from four community-based group family medicine clinics indicated that their physicians had advised them to limit or reduce the amount of fat in their diets.¹⁹

Effectiveness of Dietary Counseling

The ideal evidence to support behavioral dietary counseling would link counseling directly to improved health outcomes in randomized controlled clinical trials. In the absence of such evidence, the clinical logic behind counseling is based on a chain of critical assumptions: (1) the clinician must be able to assess whether a patient is consuming a healthy diet, (2) critical components of counseling must be routinely replicable, (3) counseling must lead to sustained improvements in diet, and (4) the health benefits of these changes in diet must be established and known to exceed the potential harms of intervention.¹¹ A review conducted for the USPSTF identified 21 fair to good quality randomized controlled clinical trials of dietary counseling among patients without existing diet-related chronic disease (eg, coronary heart disease or cancer). Trials had to include follow-up of at least 3 months after intervention for at least 50% of the enrolled subjects and include measures of dietary

intake. Studies that assessed only physiologic measures (eg, lipid levels, weight, or body mass index [BMI]) were not included. Additional details of the inclusion and exclusion criteria, and methods for assessing quality of studies, are described elsewhere.^{2,20}

Most of these trials focused exclusively on dietary counseling, though some targeted diet as part of a broader risk-factor modification program that also addressed smoking and sedentary lifestyle.²¹⁻²⁴ Most studies targeted reductions in total fat or saturated fat intake (n=17).^{7-9,13-15,21-33} Ten studies targeted increased fruit and vegetable intake^{8,9,12,21,25-27,32,34,35} and 7 targeted increased intake of fiber and whole grains.^{7,13,22,26,27,32,36} Most studies (n=11) focused on a single nutrient, although 10 focused on changes in 2 or more nutrients.^{7-9,13,21,22,25-27,32}

Studies were classified by intensity of the interventions evaluated, based on the number and length of counseling sessions, the magnitude and intensity of educational materials provided, and the use of supplemental interventions such as support group sessions or cooking classes. Low-intensity interventions involved 1 contact lasting less than 30 minutes. High-intensity interventions involved more than 6 contacts lasting more than 30 minutes. Medium-intensity interventions fell between low- and high-intensity.

Effects of counseling were classified as “large,” “medium,” or “small” for each component of diet measured.² With reference to these specific, defined categories, the USPSTF concluded that large effects sustained over time were likely to produce important health benefits (reductions in morbidity and mortality).³⁷⁻⁴¹ Given the large attributable risk associated with these dietary components, it is possible that medium or even small changes in diet would yield important health benefits across a large population. However, to date, there is little direct evidence about the effect of small and medium dietary changes on the future risk for coronary heart disease, making it difficult to determine with certainty whether such changes will translate into changes in the incidence of chronic disease. Better data about these linkages are needed.

Assessing Dietary Behaviors in Primary Care Patients

A number of brief, validated dietary assessment instruments can identify dietary counseling needs, guide intervention, and monitor change among adult patients in primary care and other clinical settings. Most of these instruments can be self-administered, are easily scored, have fewer than 40 items, and take 10 minutes or less to administer. However, these instruments are susceptible to bias (ie, patients report healthier diets than they actually consume); some studies indicate that under-reporting of caloric intake is common, especially among obese patients.¹⁰ When used to evaluate counseling efficacy, efforts to verify self-reported information are recommended.^{7-10,13,24,42} For children aged 9 years and older, food frequency questionnaires administered directly to children can provide a reasonably accurate picture of usual dietary patterns, with correlations with criterion measures ranging from 0.46 to 0.79.⁶ No brief valid dietary screening instruments were identified for children below the age of 9 years. The optimal interval for screening adults or children is not known.

Effectiveness of Routine Counseling in Primary Care

The USPSTF found 9 fair to good quality randomized controlled trials of behavioral dietary counseling in unselected populations in primary care settings. The majority of these interventions focused on change in more than one nutrient (ie, fat/saturated fat, fruit/vegetables, and/or fiber).^{7,9,13,25-27,32} Most of these trials combined basic nutrition education with behaviorally-oriented counseling to help patients acquire the skills, motivation, or support needed to alter their daily eating patterns and food selection and preparation practices. Duration of interventions lasted from 1 week to 1 year. No controlled trials with children or adolescents were identified.

The 9 studies varied in the amount of face-to-face counseling involved. Two studies of medium-intensity interventions evaluated multiple face-to-face sessions of behavioral dietary counseling provided in the primary care setting by a dietitian or nutritionist,

or by a primary care physician or nurse practitioner who had received brief training in dietary counseling.^{32,36} These interventions involved 2 to 3 group or individual sessions lasting 30 minutes, with follow-up visits at 1 and 3 months. Baron et al reported an 84% patient recruitment/participation rate.³⁶

Seven studies involved little or no face-to-face counseling and placed greater emphasis on patient self-help materials, manuals, and varied forms of interactive health communication. Two were studies of low-intensity interventions that combined brief (≤ 5 minutes) face-to-face counseling sessions with a primary care physician or nurse with self-help materials.^{7,13} Three others were studies of low-intensity interventions that relied either on mailed self-help materials^{25,34} or on health behavior change messages delivered via an automated computer-based voice system.²⁷ Campbell et al²⁵ found significantly greater benefits from tailored than non-tailored self-help materials; Lutz et al³⁴ did not. The remaining 2 were medium-intensity interventions that combined a computer-generated personalized letter and motivational phone call(s) from a trained health educator with a series of self-help mailings and newsletters.^{9,26} Patient recruitment and participation in this second group of studies ranged from 16%³⁴ to 80%,²⁵ with most in the 40% to 70% range.

These studies in unselected populations produced mostly small ($n=9$) and medium ($n=8$) as opposed to large ($n=3$) improvements in self-reported dietary behaviors, most of which were statistically significant. Most studies followed patients for 6 months or less post-intervention; 4 followed patients for as long as 12 months.^{9,13,32,36} Only 2 of them assessed impacts on intermediate biological endpoints (eg, serum cholesterol, weight, or BMI), of which none reported significant treatment effects.^{13,36} No studies examined adverse treatment effects.

The USPSTF also reviewed 2 additional studies that enrolled predominantly healthy premenopausal women, a large proportion of whom were overweight or obese. These studies employed high-intensity interventions involving multiple dietitian-led individual¹² or group³³

counseling sessions. One intervention extended over a 6-month period and aimed at increasing fruit and vegetable intake¹²; the other extended over a 5-year period and focused on dietary fat reduction. Both trials reported large treatment effects in self-reported dietary behavior at 6-month post-intervention follow-up, and both reported favorable changes in biological risk factors or markers. However, participants in these studies were highly selected and motivated volunteers. The USPSTF concluded that results could not be generalized to more representative primary care populations.

Effectiveness of Intensive Counseling in Patients at Risk for Chronic Disease

The USPSTF found 10 fair to good quality randomized controlled trials that tested whether medium- to high-intensity interventions delivered in primary care or other clinical settings led to improved dietary outcomes among adults who were identified as being at increased risk for diet-related chronic disease.^{8,14,15,21-24,28-31,35} For 2 of these trials, 2 research reports for each were reviewed.^{14,15,28,29} No controlled trials with children or adolescents at risk for chronic disease were identified that reported dietary outcomes.

The interventions involved a two-step assessment: screening to identify a patient's risk status using chart audit/clinical exam/laboratory testing to screen for hyperlipidemia, hypertension, family history of heart disease or breast cancer, overweight, obesity, smoking status, and sedentary lifestyle, followed by assessment of dietary practices using a variety of dietary assessment tools and protocols (eg, food frequency questionnaires, 3-4 day food records, and brief dietary assessment instruments). Hyperlipidemia was included as a risk factor in most of these studies. Four trials addressed diet along with physical activity and/or smoking.²¹⁻²⁴

Most of the trials tested multi-session group or individual counseling that combined nutrition education with behaviorally-oriented counseling. Most studies focused on reducing saturated fat and/or total fat intake; 2 of these studies also

targeted fiber or fruit and vegetable intake,^{21,22} and one focused on increasing fruit and vegetable intake only.³⁵ Most studies also reported intermediate health outcomes, such as serum lipid levels, blood pressure, weight, and/or BMI. Follow-up in most studies (n=6) was 12 months or longer, some as long as 4 to 6 years.^{21-24,28-30}

Six of the trials took place outside of primary care settings, where counseling was provided by an experienced nutritionist, dietitian, and/or health educator in 8 to 20 sessions over a period ranging from 4 months to 5 to 6 years.^{8,21,23,28,29,31,35} Four trials took place within primary care settings,^{14,15,22,24,30} where counseling was provided by specially trained primary care physicians or nurses (training ranging from 60 minutes to 3 days) in 3 to 6 special sessions supplemented by follow-up phone calls and/or newsletters, and follow-up at routine visits over a period of 4 to 18 months. In two primary care-based studies,^{14,15,30} behavioral dietary counseling for patients with hyperlipidemia was supplemented, if needed, with lipid-lowering medication and/or referral to outside counseling by a dietitian. Ockene et al¹⁵ found that implementing office-level systems supports (prompts, reminders, and counseling algorithms) significantly improved primary care provider adherence to the comprehensive dietary counseling.

In summary, interventions for patients at risk for chronic disease resulted in dietary behavior changes that were small (n=3),^{14,15,21,22} medium (n=6),^{8,21,22,24,30,35} and large (n=4),^{8,23,28,31} most of which were statistically significant. The magnitude and duration of these changes were greater with higher intensity interventions than with interventions of lower intensity. More than one-half of these studies found that self-reported dietary changes were accompanied by significant improvements in serum lipids, weight, or BMI.^{8,21,22,28-30} These findings help corroborate patients' self-reported dietary changes and confirm the overall health benefits of the observed changes in diet.

Discussion

Medium- to high-intensity behavioral interventions appear to produce consistent, sustained, and clinically important changes in dietary intake of total fat, saturated fat, fruit and vegetables, and fiber. However, these trials were generally either conducted with patients with known risk factors for diet-related chronic disease, or performed in special clinics with highly selected patients and specially trained providers. The most effective interventions generally combined education, behaviorally-oriented counseling, and patient reinforcement and follow-up. More intensive interventions, and those of longer duration, are associated with larger magnitude of benefit and more sustained changes in diet. Available studies do not, however, allow firm conclusions about the essential or most effective elements of these multi-component interventions, their relative effect on specific dietary constituents (eg, fat, fruit and vegetables, or fiber), or the relative efficacy of targeting single or multiple dietary risks or addressing diet in the context of broader lifestyle interventions. Although evidence is stronger for counseling patients who are at increased risk for chronic disease, such as those with hyperlipidemia, than for the general population of patients, it is not possible to disentangle the effects of patient risk status from the effects of intervention intensity. Adherence to these intensive interventions and the dietary changes they require may be dependent on patients' heightened perceived risk and motivation for change.

Existing trials of routine dietary interventions in unselected primary care populations have generally produced only small to medium changes in self-reported diet. Although direct comparisons cannot be made, results from medium-intensity, routine face-to-face counseling from nutritionists, dietitians, or specially trained primary care practitioners (physicians, nurses, or nurse practitioners) appear similar to those achieved through less intensive, minimal-contact interventions to supplement brief primary care provider advice/counseling. The consistently positive effects of such interventions on diet in unselected patient populations establish these interventions as highly promising as part of routine

preventive care for patients at average risk for chronic disease. The USPSTF concluded, however, that existing studies do not provide sufficient evidence to recommend these interventions for widespread use due to a number of limitations such as modest overall patient recruitment/participation rates, reliance on self-reported outcome measures, relatively short follow-up periods, uncertainty about the health effects of small and medium changes in diet, and the lack of evidence about possible adverse effects of counseling. Two studies suggest high-intensity interventions can be effective in selected patients at average risk, but the applicability of these findings and the feasibility of these interventions in primary care settings are uncertain.^{12,33}

Recommendations of Others

Dietary guidelines for the general population have been issued by the U.S. Department of Agriculture (USDA)¹⁶ and the Department of Health and Human Services; specific dietary objectives for the nation are outlined in *Healthy People 2010*.¹⁷ Guidelines from the American Heart Association (AHA) and the American Cancer Society (ACS) address diets that will lower the risk for heart disease and cancer, respectively.^{43,44} These guidelines generally agree in recommending a diet that includes a variety of fruit, vegetables, and grain products; is low in saturated fat and cholesterol and moderate in total fat; and balances calories with physical activity to maintain a healthy weight.

A variety of groups have recommended nutritional counseling or dietary advice for patients at average risk for chronic disease, including the American College of Preventive Medicine (ACPM), American Academy of Family Physicians (AAFP), American Academy of Pediatrics (AAP), and the American College of Obstetricians and Gynecologists (ACOG).⁴⁵⁻⁴⁸ These recommendations are based primarily on the benefits of a healthy diet rather than on evaluations of the efficacy of counseling. The Canadian Task Force on Preventive Health Care (CTFPHC) concluded in 1994 that there was fair evidence to provide general dietary advice

to all patients, based on a limited number of trials of counseling.⁴⁹

Recommendations on nutritional counseling for patients at risk (eg, those who have hypertension or hyperlipidemia) have been issued by the American Dietetic Association (ADA) and two panels sponsored by the National Institutes of Health (NIH) National Heart, Lung, and Blood Institute. The ADA recommends that primary care providers screen for nutrition-related illnesses, prescribe diets, provide preliminary counseling on specific nutritional needs, follow up with patients, and refer patients to appropriate dietetic professionals when necessary.⁵⁰ Similarly, The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure recommends that dietary assessments be included as part of routine medical history and that physicians counsel patients on lifestyle modifications for the prevention and treatment of high blood pressure (lose weight if overweight, limit alcohol intake, reduce sodium intake, reduce saturated fat and cholesterol intake).⁵¹ The National Cholesterol Education Program recommends that individuals with elevated levels of low density lipoprotein limit their intake of fats, particularly saturated fats, and cholesterol and increase dietary fiber.⁵²

References

1. U.S. Preventive Services Task Force. *Guide to Clinical Preventive Services*. 2nd ed. Washington, DC: Office of Disease Prevention and Health Promotion; 1996.
2. Ammerman A, Pignone M, Fernandez L, et al. *Counseling to Promote a Healthy Diet*. Systematic Evidence Review No. 18 (Prepared by Research Triangle Institute—University of North Carolina Evidence-Based Practice Center under Contract No. 290-97-011). Rockville, MD: Agency for Healthcare Research and Quality. April 2002. Available on the AHRQ Web site at: www.ahrq.gov/clinic/serfiles.htm.
3. Obarzanek E, Hunsberger SA, Van Horn L, et al. Safety of a fat-reduced diet: the Dietary Intervention Study in Children (DISC). *Pediatrics*. 1997;100(1):51-59.
4. Obarzanek E, Kimm SY, Barton BA, et al. Long-term safety and efficacy of a cholesterol-lowering

- diet in children with elevated low-density lipoprotein cholesterol: seven-year results of the Dietary Intervention Study in Children (DISC). *Pediatrics*. 2001;107(2):256-264.
5. Calfas KJ, Zabinski MF, Rupp J. Practical nutrition assessment in primary care settings: a review. *Am J Prev Med*. 2000;18(4):289-299.
 6. Rockett HR, Colditz GA. Assessing diets of children and adolescents. *Am J Clin Nutr*. 1997;65(4):1116-1122.
 7. Beresford SA, Farmer EM, Feingold L, Graves KL, Sumner SK, Baker RM. Evaluation of a self-help dietary intervention in a primary care setting. *Am J Public Health*. 1992;82(1):79-84.
 8. Coates RJ, Bowen DJ, Kristal AR, et al. The Women's Health Trial Feasibility Study in Minority Populations: changes in dietary intakes. *Am J Epidemiol*. 1999;149(12):1104-1112.
 9. Kristal AR, Curry SJ, Shattuck AL, Feng Z, Li S. A randomized trial of a tailored, self-help dietary intervention: the Puget Sound Eating Patterns study. *Prev Med*. 2000;31(4):380-389.
 10. Little P, Barnett J, Margetts B, et al. The validity of dietary assessment in general practice. *J Epidemiol Commun Health*. 1999;53(3):165-172.
 11. Whitlock EP, Orleans CT, Pender N, Allan J. Evaluating primary care behavioral counseling interventions: an evidence-based approach. *Am J Prev Med*. 2002;22(4):267-284.
 12. Maskarinec G, Chan CL, Meng L, Franke AA, Cooney RV. Exploring the feasibility and effects of a high-fruit and -vegetable diet in healthy women. *Cancer Epidemiol Biomarkers Prev*. 1999;8(10):919-924.
 13. Beresford SA, Curry SJ, Kristal AR, Lazovich D, Feng Z, Wagner EH. A dietary intervention in primary care practice: the Eating Patterns Study. *Am J Public Health*. 1997;87(4):610-616.
 14. Ockene IS, Hebert JR, Ockene JK, et al. Effect of physician-delivered nutrition counseling training and an office-support program on saturated fat intake, weight, and serum lipid measurements in a hyperlipidemic population: Worcester Area Trial for Counseling in Hyperlipidemia (WATCH). *Arch Int Med*. 1999;159(7):725-731.
 15. Ockene IS, Hebert JR, Ockene JK, Merriam PA, Hurley TG, Saperia GM. Effect of training and a structured office practice on physician-delivered nutrition counseling: the Worcester-Area Trial for Counseling in Hyperlipidemia (WATCH). *Am J Prev Med*. 1996;12(4):252-258.
 16. U.S. Department of Health and Human Services. Nutrition and Your Health: Dietary Guidelines for Americans, Fifth Edition (2000). Available at <http://www.health.gov/dietaryguidelines/>. Accessed November 15, 2002.
 17. U.S. Department of Health and Human Services. *Healthy People 2010: Understanding and Improving Health*. 2nd ed. Washington, DC: U.S. Government Printing Office; 2000. Available online at: <http://www.health.gov/healthypeople>.
 18. Glasgow RE, Eakin EG, Fisher EB, Bacak SJ, Brownson RC. Physician advice and support for physical activity: results from a national survey. *Am J Prev Med*. 2001;21(3):189-196.
 19. Kreuter MW, Chheda SG, Bull FC. How does physician advice influence patient behaviours? *Arch Fam Med*. 2000;9:426-433.
 20. Harris R, Helfand M, Woolf S, et al. Current methods of the U.S. Preventive Services Task Force: a review of the process. *Am J Prev Med*. 2001;20(suppl 3):21-35.
 21. Knutsen SF, Knutsen R. The Tromso Survey: the Family Intervention study—the effect of intervention on some coronary risk factors and dietary habits, a 6-year follow-up. *Prev Med*. 1991;20(2):197-212.
 22. Lindholm LH, Ekblom T, Dash C, Eriksson M, Tibblin G, Schersten B. The impact of health care advice given in primary care on cardiovascular risk. CELL Study Group. *BMJ*. 1995;310(6987):1105-1109.
 23. Neaton JD, Broste S, Cohen L, Fishman EL, Kjelsberg MO, Schoenberger J. The multiple risk factor intervention trial (MRFIT). VII. A comparison of risk factor changes between the two study groups. *Prev Med*. 1981;10(4):519-543.
 24. Steptoe A, Doherty S, Rink E, Kerry S, Kendrick T, Hilton S. Behavioural counselling in general practice for the promotion of healthy behaviour among adults at increased risk of coronary heart disease:

- randomised trial. *BMJ*. 1999;319(7215):943-947; discussion 947-948.
25. Campbell MK, DeVellis BM, Strecher VJ, Ammerman AS, DeVellis RF, Sandler RS. Improving dietary behavior: the effectiveness of tailored messages in primary care settings. *Am J Public Health*. 1994;84(5):783-787.
 26. Delichatsios HK, Hunt MK, Lobb R, Emmons K, Gillman MW. EatSmart: efficacy of a multifaceted preventive nutrition intervention in clinical practice. *Prev Med*. 2001;33(2 Pt 1):91-98.
 27. Delichatsios HK, Friedman RH, Glanz K, et al. Randomized trial of a 'talking computer' to improve adults' eating habits. *Am J Health Promot*. 2001;15(4):215-224.
 28. Henderson MM, Kushi LH, Thompson DJ, et al. Feasibility of a randomized trial of a low-fat diet for the prevention of breast cancer: dietary compliance in the Women's Health Trial Vanguard Study. *Prev Med*. 1990;19(2):115-133.
 29. Insull WJ, Henderson MM, Prentice RL, et al. Results of a randomized feasibility study of a low-fat diet. *Arch Intern Med*. 1990;150(2):421-427.
 30. Keyserling TC, Ammerman AS, Davis CE, Mok MC, Garrett J, Simpson RJ. A randomized controlled trial of a physician-directed treatment program for low-income patients with high blood cholesterol: the Southeast Cholesterol Project. *Arch Fam Med*. 1997;6(2):135-145.
 31. Mojonier ML, Hall Y, Berkson DM, et al. Experience in changing food habits of hyperlipidemic men and women. *J Am Diet Assoc*. 1980;77(2):140-148.
 32. Roderick P, Ruddock V, Hunt P, Miller G. A randomized trial to evaluate the effectiveness of dietary advice by practice nurses in lowering diet-related coronary heart disease risk. *Br J Gen Pract*. 1997;47(414):7-12.
 33. Simkin-Silverman LR, Wing RR. Management of obesity in primary care. *Obesity Research*. 1997;5(6):603-612.
 34. Lutz SF, Ammerman AS, Atwood JR, Campbell MK, DeVellis RF, Rosamond WD. Innovative newsletter interventions improve fruit and vegetable consumption in healthy adults. *J Am Diet Assoc*. 1999;99(6):705-709.
 35. Siero FW, Broer J, Bemelmans WJ, Meyboom-de Jong BM. Impact of group nutrition education and surplus value of Prochaska-based stage-matched information on health-related cognitions and on Mediterranean nutrition behavior. *Health Educ Res*. 2000;15(5):635-647.
 36. Baron JA, Gleason R, Crowe B, Mann JJ. Preliminary trial of the effect of general practice based nutritional advice. *Br J Gen Pract*. 1990;40(333):137-141.
 37. Ascherio A, Hennekens C, Willett WC, et al. Prospective study of nutritional factors, blood pressure, and hypertension among US women. *Hypertension*. 1996;27(5):1065-1072.
 38. Bazzano LA, He J, Ogden LG, et al. Legume consumption and risk of coronary heart disease in US men and women: NHANES I Epidemiologic Follow-up Study. *Arch Intern Med*. 2001;161(21):2573-2578.
 39. Jacobs DR, Pereira MA, Meyer KA, Kushi LH. Fiber from whole grains, but not refined grains, is inversely associated with all-cause mortality in older women: the Iowa women's health study. *J Am Coll Nutr*. 2000;19(suppl 3):326S-330S.
 40. Joshipura KJ, Hu FB, Manson JE, et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Intern Med*. 2001;134(12):1106-1114.
 41. Liu S, Manson JE, Stampfer MJ, et al. Whole grain consumption and risk of ischemic stroke in women: A prospective study. *JAMA*. 2000;284(12):1534-1540.
 42. Stevens VJ, Glasgow RE, Toobert DJ, Karanja N, Smith KS. Randomized trial of a brief dietary intervention to decrease consumption of fat and increase consumption of fruits and vegetables. *Am J Health Promot*. 2002;16(3):129-134.
 43. Wylie-Rosett J. Fat substitutes and health: an advisory from the Nutrition Committee of the American Heart Association. *Circulation*. 2002;105:2800-2804.
 44. American Cancer Society. Recommendations for nutrition and physical activity for cancer prevention. Available at: <http://www.cancer.org>. Accessed August 28, 2002.
 45. Nawaz H, Katz DL. American College of Preventive Medicine Policy Statement: weight management

- counseling of overweight adults. *Am J Prev Med.* 2001;21(1):73-78.
46. American Academy of Family Physicians. Summary of policy recommendations for periodic health examinations, revision 5.1. Available at: <http://www.aafp.org/exam.xml>. Accessed August 28, 2002.
 47. American Academy Of Pediatrics. Committee on Nutrition. Policy statement: Cholesterol in Childhood (RE9805). *Pediatrics.* 1998;101(1):141-147. Available at: <http://www.aap.org/policy/re9805.html>. Accessed August 28, 2002.
 48. American College of Obstetricians and Gynecologists. *Guidelines for Women's Health Care.* 2nd ed. Washington, DC: ACOG; 2002:121-34, 196-200.
 49. Canadian Task Force on the Periodic Health Examination. Canadian Guide to Clinical Preventive Health Care. Ottawa: Health Canada; 1994:586-599. Available at: <http://www.ctfphc.org>. Accessed August 28, 2002.
 50. Maillet JO, Young EA. Nutrition education for healthcare professionals: Position of the ADA. *J Am Diet Assoc.* 1998;98:343-346.
 51. Sixth report of the Joint National Committee on Prevention, Detection, Evaluation and Treatment of High Blood Pressure. Bethesda, MD: National Heart Lung and Blood Institute; 1997. Available at: <http://www.nhlbi.nih.gov/guidelines/hypertension/jncintro.htm>. Accessed August 28, 2002.
 52. Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Bethesda, MD: National Heart Lung and Blood Institute; 2001. Available at: <http://www.nhlbi.nih.gov/guidelines/cholesterol/index.htm>. Accessed August 28, 2002.

Appendix A U.S. Preventive Services Task Force—Recommendations and Ratings

The Task Force grades its recommendations according to one of 5 classifications (A, B, C, D, I) reflecting the strength of evidence and magnitude of net benefit (benefits minus harms):

- A.** The USPSTF strongly recommends that clinicians routinely provide [the service] to eligible patients. *The USPSTF found good evidence that [the service] improves important health outcomes and concludes that benefits substantially outweigh harms.*
- B.** The USPSTF recommends that clinicians routinely provide [the service] to eligible patients. *The USPSTF found at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms.*
- C.** The USPSTF makes no recommendation for or against routine provision of [the service]. *The USPSTF found at least fair evidence that [the service] can improve health outcomes but concludes that the balance of benefits and harms is too close to justify a general recommendation.*
- D.** The USPSTF recommends against routinely providing [the service] to asymptomatic patients. *The USPSTF found at least fair evidence that [the service] is ineffective or that harms outweigh benefits.*
 - I.** The USPSTF concludes that the evidence is insufficient to recommend for or against routinely providing [the service]. *Evidence that [the service] is effective is lacking, of poor quality, or conflicting and the balance of benefits and harms cannot be determined.*

Appendix B U.S. Preventive Services Task Force—Strength of Overall Evidence

The USPSTF grades the quality of the overall evidence for a service on a 3-point scale (good, fair, poor):

- Good:** Evidence includes consistent results from well-designed, well-conducted studies in representative populations that directly assess effects on health outcomes.
- Fair:** Evidence is sufficient to determine effects on health outcomes, but the strength of the evidence is limited by the number, quality, or consistency of the individual studies, generalizability to routine practice, or indirect nature of the evidence on health outcomes.
- Poor:** Evidence is insufficient to assess the effects on health outcomes because of limited number or power of studies, important flaws in their design or conduct, gaps in the chain of evidence, or lack of information on important health outcomes.

Members of the U.S. Preventive Services Task Force

Alfred O. Berg, MD, MPH, Chair, USPSTF (Professor and Chair, Department of Family Medicine, University of Washington, Seattle, WA)

Janet D. Allan, PhD, RN, Vice-chair, USPSTF (Dean, School of Nursing, University of Maryland—Baltimore, Baltimore, MD)

Paul Frame, MD (Tri-County Family Medicine, Cohocton, NY, and Clinical Professor of Family Medicine, University of Rochester, Rochester, NY)

Charles J. Homer, MD, MPH* (Executive Director, National Initiative for Children's Healthcare Quality, Boston, MA)

Mark S. Johnson, MD, MPH (Chair, Department of Family Medicine, University of Medicine and Dentistry of New Jersey—New Jersey Medical School, Newark, NJ)

Jonathan D. Klein, MD, MPH (Associate Professor, Department of Pediatrics, University of Rochester School of Medicine, Rochester, NY)

Tracy A. Lieu, MD, MPH* (Associate Professor, Department of Ambulatory Care and Prevention, Harvard Pilgrim Health Care and Harvard Medical School, Boston, MA)

Cynthia D. Mulrow, MD, MSc* (Clinical Professor and Director, Department of Medicine, University of Texas Health Science Center, San Antonio, TX)

Tracy C. Orleans, PhD (Senior Scientist and Senior Program Officer, The Robert Wood Johnson Foundation, Princeton, NJ)

Jeffrey F. Peipert, MD, MPH* (Director of Research, Women and Infants' Hospital, Providence, RI)

Nola J. Pender, PhD, RN* (Professor Emeritus, University of Michigan, Ann Arbor, MI)

Albert L. Siu, MD, MSPH (Professor of Medicine, Chief of Division of General Internal Medicine, Mount Sinai School of Medicine, New York, NY)

Steven M. Teutsch, MD, MPH (Senior Director, Outcomes Research and Management, Merck & Company, Inc., West Point, PA)

Carolyn Westhoff, MD, MSc (Professor of Obstetrics and Gynecology and Professor of Public Health, Columbia University, New York, NY)

Steven H. Woolf, MD, MPH (Professor, Department of Family Practice and Department of Preventive and Community Medicine, Virginia Commonwealth University, Fairfax, VA).

* Member of the USPSTF at the time this recommendation was finalized.



