

▼ Zyban (bupropion hydrochloride) - Safety update

Remember Zyban is contraindicated in patients with previous or current seizure disorder

Zyban was licensed in June 2000 as an aid to smoking cessation in combination with motivational support in nicotine-dependent patients. Information on the safety profile of Zyban can be found in the authorised Summary of Product Characteristics (SPC), and the Patient Information Leaflet (PIL). Up to 31 July 2001 it is estimated that 462,600 patients have received Zyban in the UK.

Up to 24 September 2001, a total of 6,570 reports of *suspected* adverse reactions have been received via the Yellow Card Scheme in the UK. The reactions most commonly reported through the Yellow Card Scheme are shown in the table below - all of these are recognised reactions and listed in the SPC and PIL. It is important to note that the *suspected* reactions are not necessarily caused by the drug and may relate to other factors such as nicotine withdrawal, other illnesses or other medicines taken concurrently.

Reported reaction	Number of reports	Reported reaction	Number of reports
Insomnia	874	Dyspnoea	214
Urticaria	859	Palpitations	214
Rashes	755	Dry mouth	213
Headache	642	Vomiting	192
Dizziness	633	Agitation	186
Nausea	559	Increased Sweating	169
Depression	426	Chest Tightness	166
Angioedema	394	Seizures	160
Tremor	316	Arthralgia	157
Pruritus	313	Constipation	152
Chest pain	293	Abdominal Pain	142
Anxiety	278	Malaise	139

(Please note that many reports contain more than one of the above-mentioned reactions, and therefore the sum of the number of reports in this table exceeds the total number of reports received for Zyban.)

Zyban is associated with a dose-related risk of seizure with an estimated incidence of approximately 0.1% (1/1,000). There have been 160 reports in the UK of seizures *suspected* as being associated with the use of Zyban. In approximately one-half of the reports, patients had either a past history of seizure(s) and/or risk factors for their occurrence.

To reduce the risk of seizures, prescribers are reminded that Zyban is contraindicated in patients with a current seizure disorder or any history of seizures, with current or previous diagnosis of bulimia or anorexia nervosa, with a known central nervous system (CNS) tumour, and those experiencing abrupt withdrawal from alcohol or benzodiazepines.

Furthermore, Zyban should not be prescribed in patients with other risk factors for seizures unless there is compelling clinical justification for which the potential benefit outweighs the increased risk of seizure. Such risk factors include concomitant use of any drug known to lower the seizure threshold (including antipsychotics, antidepressants, antimalarials, theophylline, systemic steroids, tramadol, quinolones and sedating antihistamines), alcohol abuse, a history of head trauma, diabetes treated with hypoglycaemics or insulin and use of stimulants or anorectic products. In such patients a lower dose of 150mg daily throughout the entire treatment period should be considered.

There have been 51 reports of *suspected* adverse reactions to Zyban which had a fatal outcome. The contribution of Zyban to these fatal cases is unproven and in the majority of cases the individual's underlying condition may provide an alternative explanation.

As with all new drugs, the safety of Zyban remains under close review. Doctors and pharmacists are asked to continue to report all suspected adverse reactions to the MCA/CSM.