



PDF

## Updated Information regarding Mosquito Repellents April 22, 2005

**Repellents are an important tool to assist people in protecting themselves from mosquito-borne diseases.**

A wide variety of insect repellent products are available. CDC recommends the use of products containing active ingredients which have been registered with the U.S. Environmental Protection Agency (EPA) for use as repellents applied to skin and clothing. EPA registration of repellent active ingredients indicates the materials have been reviewed and approved for efficacy and human safety when applied according to the instructions on the label.

Of the active ingredients registered with the EPA, two have demonstrated a higher degree of efficacy in the peer-reviewed, scientific literature \*. Products containing these active ingredients typically provide longer-lasting protection than others:

- DEET (N,N-diethyl-m-toluamide)
- Picaridin (KBR 3023)

Oil of lemon eucalyptus [p-menthane 3,8-diol (PMD)], a plant based repellent, is also registered with EPA. In two recent scientific publications, when oil of lemon eucalyptus was tested against mosquitoes found in the US it provided protection similar to repellents with low concentrations of DEET.

Oil of lemon eucalyptus has not been tested against mosquitoes that spread malaria and some other diseases which occur internationally. See CDC Travelers' Health website (<http://www.cdc.gov/travel/bugs.htm>) for specific recommendations concerning protection from insects when traveling outside the United States.

In addition, certain products which contain permethrin are recommended for use on clothing, shoes, bed nets, and camping gear, and are registered with EPA for this use. Permethrin is highly effective as an insecticide and as a repellent. Permethrin-treated clothing repels and kills ticks, mosquitoes, and other arthropods and retains this effect after repeated laundering. The permethrin insecticide should be reapplied following the label instructions. Some commercial products are available pretreated with permethrin.

Length of protection from mosquito bites varies with the amount of active ingredient, ambient temperature, amount of physical activity/perspiration, any water exposure, abrasive removal, and other factors. For long duration protection use a long lasting (micro-encapsulated) formula and re-apply as necessary, according to label instructions.

EPA recommends the following precautions when using insect repellents:

- Apply repellents only to exposed skin and/or clothing (as directed on the product label.) Do not use repellents under clothing.
- Never use repellents over cuts, wounds or irritated skin.

April 22, 2005

Page 1 of 2

## Updated Information regarding Mosquito Repellents

April 22, 2005

(continued from previous page)

- Do not apply to eyes or mouth, and apply sparingly around ears. When using sprays, do not spray directly on face—spray on hands first and then apply to face.
- Do not allow children to handle the product. When using on children, apply to your own hands first and then put it on the child. You may not want to apply to children's hands.
- Use just enough repellent to cover exposed skin and/or clothing. Heavy application and saturation are generally unnecessary for effectiveness. If biting insects do not respond to a thin film of repellent, then apply a bit more.
- After returning indoors, wash treated skin with soap and water or bathe. This is particularly important when repellents are used repeatedly in a day or on consecutive days. Also, wash treated clothing before wearing it again. (This precaution may vary with different repellents—check the product label.)
- If you or your child get a rash or other bad reaction from an insect repellent, stop using the repellent, wash the repellent off with mild soap and water, and call a local poison control center for further guidance. If you go to a doctor because of the repellent, take the repellent with you to show the doctor.

Note that the label for products containing oil of lemon eucalyptus specifies that they should not be used on children under the age of three years. Other than those listed above, EPA does not recommend any additional precautions for using registered repellents on pregnant or lactating women, or on children. For additional information regarding the use of repellent on children, please see CDC's Frequently Asked Questions about Repellent Use. [[http://www.cdc.gov/ncidod/dvbid/westnile/qa/insect\\_repellent.htm](http://www.cdc.gov/ncidod/dvbid/westnile/qa/insect_repellent.htm)]

DEET-based repellents applied according to label instructions may be used along with a separate sunscreen. No data are available at this time regarding the use of other active repellent ingredients in combination with a sunscreen.

See <http://www.epa.gov/pesticides/factsheets/insectrp.htm> for additional information on using EPA-registered repellents.

\*Fradin MS, Day JF. Comparative efficacy of insect repellents against mosquito bites. *N Engl J Med*. 2002; 347(1): 13-8.

Barnard DR, Xue RD. Laboratory evaluation of mosquito repellents against *Aedes albopictus*, *Culex nigripalpus*, and *Ochlerotatus triseriatus* (Diptera: Culicidae). *J Med Entomol*. 2004 Jul; 41(4): 726-30.

For more information, visit <http://www.cdc.gov/westnile>, or call CDC at 800-CDC-INFO (English and Spanish) or 888-232-6348 (TTY).